

Integrality over ideal semifiltrations

(detailed version)

Darij Grinberg

July 14, 2019*

Abstract. We study integrality over rings (all commutative in this paper) and over ideal semifiltrations (a generalization of integrality over ideals). We begin by reproving classical results, such as a version of the “faithful module” criterion for integrality over a ring, the transitivity of integrality, and the theorem that sums and products of integral elements are again integral. Then, we define the notion of integrality over an ideal semifiltration (a sequence (I_0, I_1, I_2, \dots) of ideals satisfying $I_0 = A$ and $I_a I_b \subseteq I_{a+b}$ for all $a, b \in \mathbb{N}$), which generalizes both integrality over a ring and integrality over an ideal (as considered, e.g., in Swanson/Huneke [5]). We prove a criterion that reduces this general notion to integrality over a ring using a variant of the Rees algebra. Using this criterion, we study this notion further and obtain transitivity and closedness under sums and products for it as well. Finally, we prove the curious fact that if u, x and y are three elements of a (commutative) A -algebra (for A a ring) such that u is both integral over $A[x]$ and integral over $A[y]$, then u is integral over $A[xy]$. We generalize this to integrality over ideal semifiltrations, too.

Contents

0. Definitions and notations	4
1. Integrality over rings	7
1.1. The fundamental equivalence	7
1.2. Transitivity of integrality	14

*updated and improved version of undergraduate work from 2010

1.3. Integrality of sums and products	16
1.4. Some further consequences	20
2. Integrality over ideal semifiltrations	26
2.1. Definitions of ideal semifiltrations and integrality over them	26
2.2. Polynomial rings and Rees algebras	27
2.3. Reduction to integrality over rings	30
2.4. Sums and products again	34
2.5. Transitivity again	37
3. Generalizing to two ideal semifiltrations	40
3.1. The product of two ideal semifiltrations	40
3.2. Half-reduction	41
3.3. Integrality of products over the product semifiltration	47
4. Accelerating ideal semifiltrations	49
4.1. Definition of λ -acceleration	49
4.2. Half-reduction and reduction	50
5. On a lemma by Lombardi	58
5.1. A lemma on products of powers	58
5.2. Integrality over $A[x]$ and over $A[y]$ implies integrality over $A[xy]$	69
5.3. Generalization to ideal semifiltrations	71
5.4. Second proof of Corollary 1.12	73

Introduction

The purpose of this paper is to state (and prove) some theorems and proofs related to integrality in commutative algebra in somewhat greater generality than is common in the literature. I claim no novelty, at least not for the underlying ideas, but I hope that this paper will be useful as a reference (at least for myself).

Section 1 (Integrality over rings) mainly consists of known facts (Theorem 1.1, Theorem 1.5, Theorem 1.7) and a generalized exercise from [4] (Corollary 1.12) with a few minor variations (Theorem 1.11 and Corollary 1.13).

Section 2 (Integrality over ideal semifiltrations) merges the concept of integrality over rings (as considered in Section 1) and integrality over ideals (a less popular but still highly useful notion; the book [5] is devoted to it) into one general notion: that of integrality over ideal semifiltrations (Definition 2.3). This notion is very general, yet it can be reduced to the basic notion of integrality over rings by a suitable change of base ring (Theorem 2.11). This reduction allows to extend some standard properties of integrality over rings to the general case (Theorem 2.13, Theorem 2.14 and Theorem 2.16).

Section 3 (Generalizing to two ideal semifiltrations) continues Section 2, adding

one more layer of generality. Its main results are a “relative” version of Theorem 2.11 (Theorem 3.2) and a known fact generalized once more (Theorem 3.4).

Section 4 (Accelerating ideal semifiltrations) generalizes Theorem 3.2 (and thus also Theorem 2.11) a bit further by considering accelerated ideal semifiltrations (a generalization of powers of an ideal).

Section 5 (On a lemma by Lombardi) is about an auxiliary result Henri Lombardi used in [6] to prove Kronecker’s Theorem¹. Here we show a variant of this result (generalized in one direction, less general in another).

This paper is supposed to be self-contained (only linear algebra and basic knowledge about rings, modules, ideals and polynomials is assumed).

All proofs given in this paper are constructive.

Note on the level of detail

This is the long version of this paper, with all proofs maximally detailed. For all practical purposes, the brief version [7] should be sufficient (and quite possibly easier to read).

Note on an old preprint

This is an updated and somewhat generalized version of my preprint “A few facts on integrality”, which is still available in its old form as well:

- brief version:
<https://www.cip.ifi.lmu.de/~grinberg/IntegralityBRIEF.pdf>
- long version:
<https://www.cip.ifi.lmu.de/~grinberg/Integrality.pdf> .

Be warned that said preprint has been written in 2009–2010 when I was an undergraduate, and suffers from bad writing and formatting.

Acknowledgments

I thank Irena Swanson and Marco Fontana for enlightening conversations, and Irena Swanson in particular for making her book [5] freely available (which helped me discover the subject as an undergraduate).

¹**Kronecker’s Theorem.** Let B be a ring (“ring” always means “commutative ring with unity” in this paper). Let g and h be two elements of the polynomial ring $B[X]$. Let g_α be any coefficient of the polynomial g . Let h_β be any coefficient of the polynomial h . Let A be a subring of B which contains all coefficients of the polynomial gh . Then, the element $g_\alpha h_\beta$ of B is integral over the subring A .

0. Definitions and notations

We begin our study of integrality with some classical definitions and conventions from commutative algebra:

Definition 0.1. In the following, “ring” will always mean “commutative ring with unity”. Furthermore, if A is a ring, then “ A -algebra” shall always mean “commutative A -algebra with unity”. The unity of a ring A will be denoted by 1_A or by 1 if no confusion can arise.

We denote the set $\{0, 1, 2, \dots\}$ by \mathbb{N} , and the set $\{1, 2, 3, \dots\}$ by \mathbb{N}^+ .

Definition 0.2. Let A be a ring. Let M be an A -module.

If $n \in \mathbb{N}$, and if m_1, m_2, \dots, m_n are n elements of M , then we define an A -submodule $\langle m_1, m_2, \dots, m_n \rangle_A$ of M by

$$\langle m_1, m_2, \dots, m_n \rangle_A = \left\{ \sum_{i=1}^n a_i m_i \mid (a_1, a_2, \dots, a_n) \in A^n \right\}.$$

This A -submodule $\langle m_1, m_2, \dots, m_n \rangle_A$ is known as the A -submodule of M generated by m_1, m_2, \dots, m_n (or as the A -linear span of m_1, m_2, \dots, m_n). It consists of all A -linear combinations of m_1, m_2, \dots, m_n , and in particular contains all n elements m_1, m_2, \dots, m_n . Thus, it satisfies $\{m_1, m_2, \dots, m_n\} \subseteq \langle m_1, m_2, \dots, m_n \rangle_A$.

Also, if S is a finite set, and m_s is an element of M for every $s \in S$, then we define an A -submodule $\langle m_s \mid s \in S \rangle_A$ of M by

$$\langle m_s \mid s \in S \rangle_A = \left\{ \sum_{s \in S} a_s m_s \mid (a_s)_{s \in S} \in A^S \right\}.$$

This A -submodule $\langle m_s \mid s \in S \rangle_A$ is known as the A -submodule of M generated by the family $(m_s)_{s \in S}$ (or as the A -linear span of $(m_s)_{s \in S}$). It consists of all A -linear combinations of the elements m_s with $s \in S$, and in particular contains all these elements themselves.

Of course, if m_1, m_2, \dots, m_n are n elements of M , then

$$\langle m_1, m_2, \dots, m_n \rangle_A = \langle m_s \mid s \in \{1, 2, \dots, n\} \rangle_A.$$

Let us observe a trivial fact that we shall use (often tacitly):

Lemma 0.3. Let A be a ring. Let M be an A -module. Let N be an A -submodule of M . Let S be a finite set; let m_s be an element of N for every $s \in S$. Then, $\langle m_s \mid s \in S \rangle_A \subseteq N$.

Proof of Lemma 0.3. We have $m_s \in N$ for every $s \in S$. Thus, $\sum_{s \in S} a_s m_s \in N$ for every $(a_s)_{s \in S} \in A^S$ (since N is an A -submodule of M , and thus is closed under A -linear combination). But the definition of $\langle m_s \mid s \in S \rangle_A$ yields

$$\langle m_s \mid s \in S \rangle_A = \left\{ \sum_{s \in S} a_s m_s \mid (a_s)_{s \in S} \in A^S \right\} \subseteq N$$

(since $\sum_{s \in S} a_s m_s \in N$ for every $(a_s)_{s \in S} \in A^S$). This proves Lemma 0.3. \square

Definition 0.4. Let A be a ring, and let $n \in \mathbb{N}$. Let M be an A -module. We say that the A -module M is *n-generated* if there exist n elements m_1, m_2, \dots, m_n of M such that $M = \langle m_1, m_2, \dots, m_n \rangle_A$. In other words, the A -module M is *n-generated* if and only if there exists a set S and an element m_s of M for every $s \in S$ such that $|S| = n$ and $M = \langle m_s \mid s \in S \rangle_A$.

We shall use the standard basic properties of submodules of algebras, such as the following:

Proposition 0.5. Let A be a ring. Let B be an A -algebra.

(a) For any two A -submodules U and V of B , we let $U \cdot V$ denote the A -submodule of B spanned by all products of the form uv with $(u, v) \in U \times V$. This A -submodule $U \cdot V$ is also denoted by UV . Thus we have defined a binary operation \cdot on the set of all A -submodules of B . Equipped with this operation, the set of all A -submodules of B becomes an abelian monoid, with neutral element $A \cdot 1_B$.

This all applies, in particular, to the case when $B = A$; in this case, the A -submodules of B are the same as the ideals of A . Thus, the set of all ideals of A becomes an abelian monoid, with neutral element $A \cdot 1_A = A$.

Likewise, we can define $U \cdot V$ when U is an ideal of A while V is an A -submodule of B . These “product” operations satisfy the rules one would expect, such as

$$U(V + W) = UV + UW; \quad (U + V)W = UW + VW; \quad (UV)W = U(VW)$$

(whenever these expressions make sense).

(b) Let S be a finite set. Let m_s be an element of B for each $s \in S$. Then, for any $b \in B$, we have

$$b \cdot \langle m_s \mid s \in S \rangle_A = \langle bm_s \mid s \in S \rangle_A.$$

(c) Let S be a finite set. Let m_s be an element of B for each $s \in S$. Let T be a finite set. Let n_t be an element of B for each $t \in T$. Then,

$$\langle m_s \mid s \in S \rangle_A \cdot \langle n_t \mid t \in T \rangle_A = \langle m_s n_t \mid (s, t) \in S \times T \rangle_A.$$

Definition 0.6. Let A be a ring. Let B be an A -algebra. (Let us recall that both rings and algebras are always understood to be commutative and unital in this paper.)

If u_1, u_2, \dots, u_n are n elements of B , then we define an A -subalgebra $A[u_1, u_2, \dots, u_n]$ of B by

$$A[u_1, u_2, \dots, u_n] = \{P(u_1, u_2, \dots, u_n) \mid P \in A[X_1, X_2, \dots, X_n]\}$$

(where $A[X_1, X_2, \dots, X_n]$ denotes the polynomial ring in n indeterminates X_1, X_2, \dots, X_n over A).

In particular, if u is an element of B , then the A -subalgebra $A[u]$ of B is defined by

$$A[u] = \{P(u) \mid P \in A[X]\}$$

(where $A[X]$ denotes the polynomial ring in a single indeterminate X over A). Since

$$A[X] = \left\{ \sum_{i=0}^m a_i X^i \mid m \in \mathbb{N} \text{ and } (a_0, a_1, \dots, a_m) \in A^{m+1} \right\},$$

this becomes

$$\begin{aligned} A[u] &= \left\{ \left(\sum_{i=0}^m a_i X^i \right) (u) \mid m \in \mathbb{N} \text{ and } (a_0, a_1, \dots, a_m) \in A^{m+1} \right\} \\ &\quad \left(\begin{array}{l} \text{where } \left(\sum_{i=0}^m a_i X^i \right) (u) \text{ means the} \\ \text{polynomial } \sum_{i=0}^m a_i X^i \text{ evaluated at } X = u \end{array} \right) \\ &= \left\{ \sum_{i=0}^m a_i u^i \mid m \in \mathbb{N} \text{ and } (a_0, a_1, \dots, a_m) \in A^{m+1} \right\} \\ &\quad \left(\text{because } \left(\sum_{i=0}^m a_i X^i \right) (u) = \sum_{i=0}^m a_i u^i \right). \end{aligned}$$

Obviously, $uA[u] \subseteq A[u]$ (since $A[u]$ is an A -algebra and $u \in A[u]$).

Definition 0.7. Let B be a ring, and let A be a subring of B . Then, B canonically becomes an A -algebra. The A -module structure of this A -algebra B is given by multiplication inside B .

Definition 0.7 shows that theorems about A -algebras (for a ring A) are always more general than theorems about rings that contain A as a subring. Hence, we shall study A -algebras in the following, even though most of the applications of the results we shall see are found at the level of rings containing A .

1. Integrality over rings

1.1. The fundamental equivalence

Most of the theory of integrality is based upon the following result:

Theorem 1.1. Let A be a ring. Let B be an A -algebra. Thus, B is canonically an A -module. Let $n \in \mathbb{N}$. Let $u \in B$. Then, the following four assertions \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} are equivalent:

- *Assertion \mathcal{A} :* There exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$.
- *Assertion \mathcal{B} :* There exist a B -module C and an n -generated A -submodule U of C such that $uU \subseteq U$ and such that every $v \in B$ satisfying $vU = 0$ satisfies $v = 0$. (Here, C is an A -module, since C is a B -module and B is an A -algebra.)
- *Assertion \mathcal{C} :* There exists an n -generated A -submodule U of B such that $1 \in U$ and $uU \subseteq U$. (Here and in the following, “1” means “ 1_B ”, that is, the unity of the ring B .)
- *Assertion \mathcal{D} :* We have $A[u] = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$.

We shall soon prove this theorem; first, let us explain what it is for:

Definition 1.2. Let A be a ring. Let B be an A -algebra. Let $n \in \mathbb{N}$. Let $u \in B$. We say that the element u of B is *n -integral over A* if it satisfies the four equivalent assertions \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} of Theorem 1.1.

Hence, in particular, the element u of B is *n -integral over A* if and only if it satisfies the assertion \mathcal{A} of Theorem 1.1. In other words, u is *n -integral over A* if and only if there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$.

The notion of “ n -integral” elements that we have just defined is a refinement of the classical notion of integrality of elements over rings (see, e.g., [1, Definition (10.21)] or [2, Chapter V, §1.1, Definition 1] or [3, Definition 8.1.1] for this classical notion, and [5, Definition 2.1.1] for its particular case when A is a subring of B). Indeed, the classical notion defines an element u of B to be *integral over A* if and only if (using the language of our Definition 1.2) there exists some $n \in \mathbb{N}$ such that u is *n -integral over A* . Since I believe the concrete value of n to be worth more than its mere existence, I prefer the specificity of the “ n -integral” concept to the slickness of “integral”.

Theorem 1.1 is one of several similar results providing equivalent criteria for the integrality of an element of an A -algebra. See [1, Proposition (10.23)], [2,

Chapter V, Section 1.1, Theorem 1] or [3, Theorem 8.1.6] for other such results (some very close to Theorem 1.1, and all proven in similar ways).

Before we prove Theorem 1.1, let us recall a classical property of matrices:

Lemma 1.3. Let B be a ring. Let $n \in \mathbb{N}$. Let M be an $n \times n$ -matrix over B . Then,

$$\det M \cdot I_n = \operatorname{adj} M \cdot M.$$

(Here, I_n means the $n \times n$ identity matrix and $\operatorname{adj} M$ denotes the adjugate of the matrix M . The expressions “ $\det M \cdot I_n$ ” and “ $\operatorname{adj} M \cdot M$ ” have to be understood as “ $(\det M) \cdot I_n$ ” and “ $(\operatorname{adj} M) \cdot M$ ”, respectively.)

Lemma 1.3 is well-known (for example, it follows from [8, Theorem 6.100], applied to $\mathbb{K} = B$ and $A = M$).

Proof of Theorem 1.1. We will prove the implications $\mathcal{A} \implies \mathcal{C}$, $\mathcal{C} \implies \mathcal{B}$, $\mathcal{B} \implies \mathcal{A}$, $\mathcal{A} \implies \mathcal{D}$ and $\mathcal{D} \implies \mathcal{C}$.

Proof of the implication $\mathcal{A} \implies \mathcal{C}$. Assume that Assertion \mathcal{A} holds. Then, there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$. Consider this P . Since $P \in A[X]$ is a monic polynomial with $\deg P = n$, there exist elements a_0, a_1, \dots, a_{n-1} of A such that $P(X) = X^n + \sum_{k=0}^{n-1} a_k X^k$. Consider these

a_0, a_1, \dots, a_{n-1} . Substituting u for X in the equality $P(X) = X^n + \sum_{k=0}^{n-1} a_k X^k$, we find $P(u) = u^n + \sum_{k=0}^{n-1} a_k u^k$. Hence, the equality $P(u) = 0$ (which holds by

definition of P) rewrites as $u^n + \sum_{k=0}^{n-1} a_k u^k = 0$. Hence, $u^n = -\sum_{k=0}^{n-1} a_k u^k$.

Let U be the A -submodule $\langle u^0, u^1, \dots, u^{n-1} \rangle_A$ of B . Then, $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$ and

$$u^n = -\sum_{k=0}^{n-1} a_k u^k \in \langle u^0, u^1, \dots, u^{n-1} \rangle_A = U.$$

Moreover, the n elements u^0, u^1, \dots, u^{n-1} belong to U (since $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$). In other words,

$$u^i \in U \quad \text{for each } i \in \{0, 1, \dots, n-1\}. \quad (1)$$

This relation also holds for $i = n$ (since $u^n \in U$); thus, it holds for all $i \in \{0, 1, \dots, n\}$. In other words, we have

$$u^i \in U \quad \text{for each } i \in \{0, 1, \dots, n\}. \quad (2)$$

Applying this to $i = 0$, we find $u^0 \in U$ (since $0 \in \{0, 1, \dots, n\}$). This rewrites as $1 \in U$ (since $u^0 = 1$).

Recall that $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$. Hence, U is an n -generated A -module (since u^0, u^1, \dots, u^{n-1} are n elements of U).

Now, for each $s \in \{0, 1, \dots, n-1\}$, we have $s+1 \in \{1, 2, \dots, n\} \subseteq \{0, 1, \dots, n\}$ and thus $u^{s+1} \in U$ (by (2), applied to $i = s+1$). Hence, Lemma 0.3 (applied to $M = B, N = U, S = \{0, 1, \dots, n-1\}$ and $m_s = u^{s+1}$) yields

$$\langle u^{s+1} \mid s \in \{0, 1, \dots, n-1\} \rangle_A \subseteq U.$$

Now, from $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$, we obtain

$$\begin{aligned} uU &= u \langle u^0, u^1, \dots, u^{n-1} \rangle_A = \langle u \cdot u^0, u \cdot u^1, \dots, u \cdot u^{n-1} \rangle_A \\ &= \left\langle \underbrace{u \cdot u^s}_{=u^{s+1}} \mid s \in \{0, 1, \dots, n-1\} \right\rangle_A = \langle u^{s+1} \mid s \in \{0, 1, \dots, n-1\} \rangle_A \subseteq U. \end{aligned}$$

Thus, we have found an n -generated A -submodule U of B such that $1 \in U$ and $uU \subseteq U$. Hence, Assertion \mathcal{C} holds. Hence, we have proved that $\mathcal{A} \implies \mathcal{C}$.

Proof of the implication $\mathcal{C} \implies \mathcal{B}$. Assume that Assertion \mathcal{C} holds. Then, there exists an n -generated A -submodule U of B such that $1 \in U$ and $uU \subseteq U$. Consider this U . Every $v \in B$ satisfying $vU = 0$ satisfies $v = 0$ (since $1 \in U$ and $vU = 0$ yield $v \cdot \underbrace{1}_{\in U} \in vU = 0$ and thus $v \cdot 1 = 0$, so that $v = 0$). Set $C = B$.

Then, C is a B -module, and U is an n -generated A -submodule of C (since U is an n -generated A -submodule of B , and $C = B$) such that $uU \subseteq U$ and such that every $v \in B$ satisfying $vU = 0$ satisfies $v = 0$. Thus, Assertion \mathcal{B} holds. Hence, we have proved that $\mathcal{C} \implies \mathcal{B}$.

Proof of the implication $\mathcal{B} \implies \mathcal{A}$. Assume that Assertion \mathcal{B} holds. Then, there exist a B -module C and an n -generated A -submodule² U of C such that $uU \subseteq U$, and such that every $v \in B$ satisfying $vU = 0$ satisfies $v = 0$. Consider these C and U .

The A -module U is n -generated. In other words, there exist n elements m_1, m_2, \dots, m_n of U such that $U = \langle m_1, m_2, \dots, m_n \rangle_A$. Consider these m_1, m_2, \dots, m_n . For any $k \in \{1, 2, \dots, n\}$, we have $m_k \in U$ (since $U = \langle m_1, m_2, \dots, m_n \rangle_A$) and thus

$$um_k \in uU \subseteq U = \langle m_1, m_2, \dots, m_n \rangle_A,$$

so that there exist n elements $a_{k,1}, a_{k,2}, \dots, a_{k,n}$ of A such that

$$um_k = \sum_{i=1}^n a_{k,i} m_i. \quad (3)$$

Consider these $a_{k,1}, a_{k,2}, \dots, a_{k,n}$.

²where C is an A -module, since C is a B -module and B is an A -algebra

The A -algebra B gives rise to a canonical ring homomorphism $\iota : A \rightarrow B$ (sending each $a \in A$ to $a \cdot 1_B \in B$). This ring homomorphism, in turn, induces a ring homomorphism $\iota^{n \times n} : A^{n \times n} \rightarrow B^{n \times n}$ (which acts on an $n \times n$ -matrix by applying ι to each entry of the matrix).

We introduce two notations:

- For any matrix T and any integers x and y , we denote by $T_{x,y}$ the entry of the matrix T in the x -th row and the y -th column.
- For any assertion \mathcal{U} , we denote by $[\mathcal{U}]$ the Boolean value of the assertion \mathcal{U} (that is, $[\mathcal{U}] = \begin{cases} 1, & \text{if } \mathcal{U} \text{ is true;} \\ 0, & \text{if } \mathcal{U} \text{ is false} \end{cases}$). This value $[\mathcal{U}]$ is an element of $\{0, 1\}$ and is also known as the *truth value* of \mathcal{U} .

Clearly, the $n \times n$ identity matrix I_n satisfies

$$(I_n)_{k,i} = [k = i] \quad \text{for every } k \in \{1, 2, \dots, n\} \text{ and } i \in \{1, 2, \dots, n\}.$$

Note that for every $k \in \{1, 2, \dots, n\}$, we have

$$m_k = \sum_{i=1}^n (I_n)_{k,i} m_i, \quad (4)$$

since

$$\begin{aligned} & \underbrace{\sum_{i=1}^n}_{=\sum_{i \in \{1, 2, \dots, n\}}} \underbrace{(I_n)_{k,i}}_{\substack{=[k=i] \\ =[i=k]}} m_i = \sum_{i \in \{1, 2, \dots, n\}} [i = k] m_i \\ &= \sum_{\substack{i \in \{1, 2, \dots, n\} \\ \text{such that } i=k}} \underbrace{[i = k]}_{=1} m_i + \sum_{\substack{i \in \{1, 2, \dots, n\} \\ \text{such that } i \neq k}} \underbrace{[i = k]}_{=0} m_i \\ &= \sum_{\substack{i \in \{1, 2, \dots, n\} \\ \text{such that } i=k}} \underbrace{1 m_i}_{=m_i} + \underbrace{\sum_{\substack{i \in \{1, 2, \dots, n\} \\ \text{such that } i \neq k}} 0 m_i}_{=0} = \sum_{\substack{i \in \{1, 2, \dots, n\} \\ \text{such that } i=k}} m_i + 0 \\ &= \sum_{\substack{i \in \{1, 2, \dots, n\} \\ \text{such that } i=k}} m_i = \sum_{i \in \{k\}} m_i \\ &\quad \left(\begin{array}{l} \text{since } \{i \in \{1, 2, \dots, n\} \mid i = k\} = \{k\} \\ \text{(because } k \in \{1, 2, \dots, n\}) \end{array} \right) \\ &= m_k. \end{aligned}$$

Hence, for every $k \in \{1, 2, \dots, n\}$, we have

$$\begin{aligned} \sum_{i=1}^n \left(u (I_n)_{k,i} - a_{k,i} \right) m_i &= \sum_{i=1}^n \left(u (I_n)_{k,i} m_i - a_{k,i} m_i \right) = u \underbrace{\sum_{i=1}^n (I_n)_{k,i} m_i}_{=m_k \text{ (by (4))}} - \sum_{i=1}^n a_{k,i} m_i \\ &= um_k - \sum_{i=1}^n a_{k,i} m_i = 0 \quad (\text{by (3)}). \end{aligned} \quad (5)$$

Define a matrix $S \in A^{n \times n}$ by

$$(S_{k,i} = a_{k,i} \text{ for all } k \in \{1, 2, \dots, n\} \text{ and } i \in \{1, 2, \dots, n\}).$$

Define a matrix $T \in B^{n \times n}$ by

$$T = \text{adj}(uI_n - S).$$

Here, the “ S ” in “ $uI_n - S$ ” means not the matrix $S \in A^{n \times n}$ itself, but rather its image under the ring homomorphism $\iota^{n \times n} : A^{n \times n} \rightarrow B^{n \times n}$; thus, the matrix $uI_n - S$ is a well-defined matrix in $B^{n \times n}$.

Let $P \in A[X]$ be the characteristic polynomial of the matrix $S \in A^{n \times n}$. Then, P is monic, and $\deg P = n$. Besides, the definition of P yields $P(X) = \det(XI_n - S)$, so that $P(u) = \det(uI_n - S)$. Therefore,

$$\begin{aligned} P(u) \cdot I_n &= \det(uI_n - S) \cdot I_n = \underbrace{\text{adj}(uI_n - S)}_{=T} \cdot (uI_n - S) \\ &\quad (\text{by Lemma 1.3, applied to } M = uI_n - S) \\ &= T \cdot (uI_n - S). \end{aligned}$$

Now, for every $\tau \in \{1, 2, \dots, n\}$, we have

$$\begin{aligned}
 P(u) \cdot m_\tau &= P(u) \cdot \sum_{i=1}^n (I_n)_{\tau,i} m_i \\
 &\left(\text{since (4) (applied to } k = \tau) \text{ yields } m_\tau = \sum_{i=1}^n (I_n)_{\tau,i} m_i \right) \\
 &= \sum_{i=1}^n \underbrace{P(u) \cdot (I_n)_{\tau,i}}_{=(P(u) \cdot I_n)_{\tau,i}} m_i = \sum_{i=1}^n \left(\underbrace{P(u) \cdot I_n}_{=(T \cdot (uI_n - S))}_{\tau,i} \right) m_i \\
 &= \sum_{i=1}^n \underbrace{(T \cdot (uI_n - S))_{\tau,i}}_{=\sum_{k=1}^n T_{\tau,k} (uI_n - S)_{k,i}} m_i = \sum_{i=1}^n \sum_{k=1}^n T_{\tau,k} (uI_n - S)_{k,i} m_i \\
 &\quad \text{(by the definition of the product of two matrices)} \\
 &= \sum_{k=1}^n T_{\tau,k} \sum_{i=1}^n \underbrace{(uI_n - S)_{k,i}}_{=u(I_n)_{k,i} - S_{k,i}} m_i = \sum_{k=1}^n T_{\tau,k} \sum_{i=1}^n \left(u(I_n)_{k,i} - \underbrace{S_{k,i}}_{=a_{k,i}} \right) m_i \\
 &= \sum_{k=1}^n T_{\tau,k} \underbrace{\sum_{i=1}^n (u(I_n)_{k,i} - a_{k,i})}_{=0}_{\text{(by (5))}} m_i = 0.
 \end{aligned}$$

But from $U = \langle m_1, m_2, \dots, m_n \rangle_A$, we obtain

$$\begin{aligned}
 P(u) \cdot U &= P(u) \cdot \langle m_1, m_2, \dots, m_n \rangle_A = \langle P(u) \cdot m_1, P(u) \cdot m_2, \dots, P(u) \cdot m_n \rangle_A \\
 &= \langle 0, 0, \dots, 0 \rangle_A \quad (\text{since } P(u) \cdot m_\tau = 0 \text{ for any } \tau \in \{1, 2, \dots, n\}) \\
 &= 0.
 \end{aligned}$$

But recall that every $v \in B$ satisfying $vU = 0$ satisfies $v = 0$. Applying this to $v = P(u)$, we find $P(u) = 0$ (since $P(u) \cdot U = 0$). Thus, we have found a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$. Therefore, Assertion \mathcal{A} holds. Hence, we have proved that $\mathcal{B} \implies \mathcal{A}$.

Proof of the implication $\mathcal{A} \implies \mathcal{D}$. Assume that Assertion \mathcal{A} holds. Then, there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$. Consider this P .

Let U be the A -submodule $\langle u^0, u^1, \dots, u^{n-1} \rangle_A$ of B . As in the Proof of the implication $\mathcal{A} \implies \mathcal{C}$, we can show that U is an n -generated A -module, and that $1 \in U$ and $uU \subseteq U$.

Now, it is easy to show that

$$u^i \in U \quad \text{for any } i \in \mathbb{N}. \quad (6)$$

[Proof of (6). We will prove (6) by induction over i :

Induction base: The assertion (6) holds for $i = 0$ (since $u^0 = 1 \in U$). This completes the induction base.

Induction step: Let $\tau \in \mathbb{N}$. If the assertion (6) holds for $i = \tau$, then the assertion (6) holds for $i = \tau + 1$ (because if the assertion (6) holds for $i = \tau$, then $u^\tau \in U$, so that $u^{\tau+1} = u \cdot \underbrace{u^\tau}_{\in U} \in uU \subseteq U$, so that $u^{\tau+1} \in U$, and thus the assertion (6) holds for $i = \tau + 1$). This completes the induction step.

Hence, the induction is complete, and (6) is proven.]

But recall that U is an A -module, and therefore is closed under A -linear combination. Thus, for any $m \in \mathbb{N}$ and any $(a_0, a_1, \dots, a_m) \in A^{m+1}$, we have $\sum_{i=0}^m a_i u^i \in U$, because each $i \in \{0, 1, \dots, m\}$ satisfies $a_i \in A$ and $u^i \in U$ (by (6)).

Now, the definition of $A[u]$ yields

$$A[u] = \left\{ \sum_{i=0}^m a_i u^i \mid m \in \mathbb{N} \text{ and } (a_0, a_1, \dots, a_m) \in A^{m+1} \right\} \subseteq U$$

(since $\sum_{i=0}^m a_i u^i \in U$ for any $m \in \mathbb{N}$ and any $(a_0, a_1, \dots, a_m) \in A^{m+1}$). On the other hand, $U \subseteq A[u]$, since

$$\begin{aligned} U &= \langle u^0, u^1, \dots, u^{n-1} \rangle_A = \left\{ \sum_{i=0}^{n-1} a_i u^i \mid (a_0, a_1, \dots, a_{n-1}) \in A^n \right\} \\ &\subseteq \left\{ \sum_{i=0}^m a_i u^i \mid m \in \mathbb{N} \text{ and } (a_0, a_1, \dots, a_m) \in A^{m+1} \right\} = A[u]. \end{aligned}$$

Combining this with $A[u] \subseteq U$, we obtain $U = A[u]$. Comparing this with $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$, we obtain $A[u] = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$.

Thus, Assertion \mathcal{D} holds. Hence, we have proved that $\mathcal{A} \implies \mathcal{D}$.

Proof of the implication $\mathcal{D} \implies \mathcal{C}$. Assume that Assertion \mathcal{D} holds. Then, $A[u] = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$.

Let U be the A -submodule $\langle u^0, u^1, \dots, u^{n-1} \rangle_A$ of B . Then, u^0, u^1, \dots, u^{n-1} are n elements of U . Hence, U is an n -generated A -module (since $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$). Comparing $U = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$ with $A[u] = \langle u^0, u^1, \dots, u^{n-1} \rangle_A$, we obtain $U = A[u]$. Now, $1 = u^0 \in A[u] = U$.

Also, from $U = A[u]$, we obtain $uU = u \cdot A[u] \subseteq A[u] = U$.

Thus, we have found an n -generated A -submodule U of B such that $1 \in U$ and $uU \subseteq U$. Hence, Assertion \mathcal{C} holds. Thus, we have proved that $\mathcal{D} \implies \mathcal{C}$.

Now, we have proved the implications $\mathcal{A} \implies \mathcal{D}$, $\mathcal{D} \implies \mathcal{C}$, $\mathcal{C} \implies \mathcal{B}$ and $\mathcal{B} \implies \mathcal{A}$ above. Thus, all four assertions \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} are equivalent, and Theorem 1.1 is proven. \square

For the sake of completeness (and as a very easy exercise), let us state a basic property of integrality that we will not ever use:

Proposition 1.4. Let A be a ring. Let B be an A -algebra. Let $u \in B$. Let $q \in \mathbb{N}$ and $p \in \mathbb{N}$ be such that $p \geq q$. Assume that u is q -integral over A . Then, u is p -integral over A .

Proof of Proposition 1.4. The element u is q -integral over A . Thus, it satisfies the Assertion \mathcal{D} of Theorem 1.1, stated for q in lieu of n . In other words, it satisfies $A[u] = \langle u^0, u^1, \dots, u^{q-1} \rangle_A$. Note that $\langle u^0, u^1, \dots, u^{p-1} \rangle_A$ is an A -submodule of B .

But $p \geq q$, thus $q \leq p$ and therefore $q - 1 \leq p - 1$. Every $s \in \{0, 1, \dots, q - 1\}$ satisfies $s \in \{0, 1, \dots, q - 1\} \subseteq \{0, 1, \dots, p - 1\}$ (since $q - 1 \leq p - 1$) and therefore $u^s \in \{u^0, u^1, \dots, u^{p-1}\} \subseteq \langle u^0, u^1, \dots, u^{p-1} \rangle_A$. Thus, u^s is an element of $\langle u^0, u^1, \dots, u^{p-1} \rangle_A$ for every $s \in \{0, 1, \dots, q - 1\}$. Hence, Lemma 0.3 (applied to $M = B$, $N = \langle u^0, u^1, \dots, u^{p-1} \rangle_A$, $S = \{0, 1, \dots, q - 1\}$ and $m_s = u^s$) shows that

$$\langle u^s \mid s \in \{0, 1, \dots, q - 1\} \rangle_A \subseteq \langle u^0, u^1, \dots, u^{p-1} \rangle_A.$$

Now,

$$A[u] = \langle u^0, u^1, \dots, u^{q-1} \rangle_A = \langle u^s \mid s \in \{0, 1, \dots, q - 1\} \rangle_A \subseteq \langle u^0, u^1, \dots, u^{p-1} \rangle_A.$$

Combining this with $\langle u^0, u^1, \dots, u^{p-1} \rangle_A \subseteq A[u]$ (which is obvious, since every A -linear combination of u^0, u^1, \dots, u^{p-1} is a polynomial in u with coefficients in A), we obtain $A[u] = \langle u^0, u^1, \dots, u^{p-1} \rangle_A$. In other words, u satisfies the Assertion \mathcal{D} of Theorem 1.1, stated for p in lieu of n . Hence, u is p -integral over A . This proves Proposition 1.4. \square

1.2. Transitivity of integrality

Let us now prove the first and probably most important consequence of Theorem 1.1:

Theorem 1.5. Let A be a ring. Let B be an A -algebra. Let $v \in B$ and $u \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Assume that v is m -integral over A , and that u is n -integral over $A[v]$. Then, u is nm -integral over A .

(Here, we are using the fact that if A is a ring, and if v is an element of an A -algebra B , then $A[v]$ is a subring of B , and therefore B is an $A[v]$ -algebra.)

Proof of Theorem 1.5. Since v is m -integral over A , we have $A[v] = \langle v^0, v^1, \dots, v^{m-1} \rangle_A$ (this is the Assertion \mathcal{D} of Theorem 1.1, stated for v and m in lieu of u and n).

Since u is n -integral over $A[v]$, we have $(A[v])[u] = \langle u^0, u^1, \dots, u^{n-1} \rangle_{A[v]}$ (this is the Assertion \mathcal{D} of Theorem 1.1, stated for $A[v]$ in lieu of A).

Let $S = \{0, 1, \dots, n-1\} \times \{0, 1, \dots, m-1\}$. Then, S is a finite set with size $|S| = nm$.

[Proof: From $S = \{0, 1, \dots, n-1\} \times \{0, 1, \dots, m-1\}$, we obtain

$$\begin{aligned} |S| &= |\{0, 1, \dots, n-1\} \times \{0, 1, \dots, m-1\}| \\ &= \underbrace{|\{0, 1, \dots, n-1\}|}_{=n} \cdot \underbrace{|\{0, 1, \dots, m-1\}|}_{=m} = nm. \end{aligned}$$

Thus, S is finite.]

Let $x \in (A[v])[u]$. Then, there exist n elements b_0, b_1, \dots, b_{n-1} of $A[v]$ such that $x = \sum_{i=0}^{n-1} b_i u^i$ (since $x \in (A[v])[u] = \langle u^0, u^1, \dots, u^{n-1} \rangle_{A[v]}$). Consider these b_0, b_1, \dots, b_{n-1} .

For each $i \in \{0, 1, \dots, n-1\}$, there exist m elements $a_{i,0}, a_{i,1}, \dots, a_{i,m-1}$ of A such that $b_i = \sum_{j=0}^{m-1} a_{i,j} v^j$ (because $b_i \in A[v] = \langle v^0, v^1, \dots, v^{m-1} \rangle_A$). Consider these $a_{i,0}, a_{i,1}, \dots, a_{i,m-1}$. Thus,

$$\begin{aligned} x &= \sum_{i=0}^{n-1} \underbrace{b_i}_{=\sum_{j=0}^{m-1} a_{i,j} v^j} u^i = \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} a_{i,j} v^j u^i = \sum_{(i,j) \in \{0,1,\dots,n-1\} \times \{0,1,\dots,m-1\}} a_{i,j} v^j u^i \\ &= \sum_{(i,j) \in S} a_{i,j} v^j u^i \quad (\text{since } \{0, 1, \dots, n-1\} \times \{0, 1, \dots, m-1\} = S) \\ &\in \langle v^j u^i \mid (i, j) \in S \rangle_A \quad (\text{since } a_{i,j} \in A \text{ for every } (i, j) \in S). \end{aligned}$$

Now, forget that we fixed x . So we have proved that $x \in \langle v^j u^i \mid (i, j) \in S \rangle_A$ for every $x \in (A[v])[u]$. In other words, $(A[v])[u] \subseteq \langle v^j u^i \mid (i, j) \in S \rangle_A$. Conversely, $\langle v^j u^i \mid (i, j) \in S \rangle_A \subseteq (A[v])[u]$ (since $v^j \in A[v]$ for every $(i, j) \in S$, and thus $\underbrace{v^j}_{\in A[v]} u^i \in (A[v])[u]$ for every $(i, j) \in S$, and therefore

$$\begin{aligned} \langle v^j u^i \mid (i, j) \in S \rangle_A &= \left\{ \begin{array}{l} \underbrace{\sum_{(i,j) \in S} a_{i,j} v^j u^i}_{\in (A[v])[u]} \quad \mid \quad (a_{i,j})_{(i,j) \in S} \in A^S \\ \text{(since } v^j u^i \in (A[v])[u] \text{ for every } (i,j) \in S \\ \text{and since } (A[v])[u] \text{ is an } A\text{-module)} \end{array} \right\} \\ &\subseteq (A[v])[u] \end{aligned}$$

). Combining these two relations, we find $(A[v])[u] = \langle v^j u^i \mid (i, j) \in S \rangle_A$. Thus, the A -module $(A[v])[u]$ is nm -generated (since $|S| = nm$).

Let $U = (A[v])[u]$. Thus, the A -module U is nm -generated (since the A -module $(A[v])[u]$ is nm -generated). Besides, U is an A -submodule of B , and we have $1 = u^0 \in (A[v])[u] = U$ and

$$\begin{aligned} uU &= u(A[v])[u] \subseteq (A[v])[u] \\ &\quad (\text{since } (A[v])[u] \text{ is an } A[v]\text{-algebra and } u \in (A[v])[u]) \\ &= U. \end{aligned}$$

Altogether, we now know that the A -submodule U of B is nm -generated and satisfies $1 \in U$ and $uU \subseteq U$.

Thus, the element u of B satisfies the Assertion \mathcal{C} of Theorem 1.1 with n replaced by nm . Hence, $u \in B$ satisfies the four equivalent assertions \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} of Theorem 1.1, all with n replaced by nm . Thus, u is nm -integral over A . This proves Theorem 1.5. \square

1.3. Integrality of sums and products

Before the next significant consequence of Theorem 1.1, let us show an essentially trivial fact:

Theorem 1.6. Let A be a ring. Let B be an A -algebra. Let $a \in A$. Then, $a \cdot 1_B \in B$ is 1-integral over A .

Proof of Theorem 1.6. The polynomial $X - a \in A[X]$ is monic and satisfies $\deg(X - a) = 1$; moreover, evaluating this polynomial at $a \cdot 1_B \in B$ yields $a \cdot 1_B - a \cdot 1_B = 0$. Hence, there exists a monic polynomial $P \in A[X]$ with $\deg P = 1$ and $P(a \cdot 1_B) = 0$ (namely, the polynomial $P \in A[X]$ defined by $P(X) = X - a$). Thus, $a \cdot 1_B$ is 1-integral over A . This proves Theorem 1.6. \square

The following theorem is a standard result, generalizing (for example) the classical fact that sums and products of algebraic integers are again algebraic integers:

Theorem 1.7. Let A be a ring. Let B be an A -algebra. Let $x \in B$ and $y \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Assume that x is m -integral over A , and that y is n -integral over A .

- (a) Then, $x + y$ is nm -integral over A .
- (b) Furthermore, xy is nm -integral over A .

Our proof of this theorem will rely on a simple lemma:

Lemma 1.8. Let A be a ring. Let C be an A -algebra. Let $x \in C$.

Let $n \in \mathbb{N}$. Let $P \in A[X]$ be a monic polynomial with $\deg P = n$. Define a polynomial $Q \in C[X]$ by $Q(X) = P(X - x)$. Then, Q is a monic polynomial with $\deg Q = n$.

Proof of Lemma 1.8. Recall that P is a monic polynomial with $\deg P = n$; hence, we can write P in the form

$$P = X^n + \sum_{i=0}^{n-1} a_i X^i \quad (7)$$

for some $a_0, a_1, \dots, a_{n-1} \in A$. Consider these a_0, a_1, \dots, a_{n-1} .

Consider the C -submodule $\langle X^0, X^1, \dots, X^{n-1} \rangle_C$ of $C[X]$. We have $X - x = X + (-x)$ and thus

$$\begin{aligned} (X - x)^n &= (X + (-x))^n = \sum_{i=0}^n \binom{n}{i} X^i (-x)^{n-i} && \text{(by the binomial formula)} \\ &= \sum_{i=0}^{n-1} \binom{n}{i} \underbrace{X^i (-x)^{n-i}}_{=(-x)^{n-i} X^i} + \binom{n}{n} X^n \underbrace{(-x)^{n-n}}_{=(-x)^0=1} \\ &\quad \text{(here, we have split off the addend for } i = n \text{ from the sum)} \\ &= \sum_{i=0}^{n-1} \binom{n}{i} (-x)^{n-i} X^i + X^n = X^n + \underbrace{\sum_{i=0}^{n-1} \binom{n}{i} (-x)^{n-i} X^i}_{\in \langle X^0, X^1, \dots, X^{n-1} \rangle_C} \\ &\in X^n + \langle X^0, X^1, \dots, X^{n-1} \rangle_C. \end{aligned} \quad (8)$$

Furthermore, for each $i \in \{0, 1, \dots, n-1\}$, we have

$$\begin{aligned} (X - x)^i &= (X + (-x))^i && \text{(since } X - x = X + (-x)) \\ &= \sum_{j=0}^i \binom{i}{j} \underbrace{X^j (-x)^{i-j}}_{=(-x)^{i-j} X^j} && \text{(by the binomial formula)} \\ &= \sum_{j=0}^i \binom{i}{j} (-x)^{i-j} X^j \in \langle X^0, X^1, \dots, X^i \rangle_C \\ &\subseteq \langle X^0, X^1, \dots, X^{n-1} \rangle_C \end{aligned} \quad (9)$$

(since $i \leq n - 1$). Now,

$$\begin{aligned}
 Q(X) &= P(X - x) = \underbrace{(X - x)^n}_{\substack{\in X^n + \langle X^0, X^1, \dots, X^{n-1} \rangle_C \\ \text{(by (8))}}} + \underbrace{\sum_{i=0}^{n-1} a_i (X - x)^i}_{\substack{\in \langle X^0, X^1, \dots, X^{n-1} \rangle_C \\ \text{(by (9))}}} \\
 &\quad \text{(here, we have substituted } X - x \text{ for } X \text{ in (7))} \\
 &\in X^n + \underbrace{\langle X^0, X^1, \dots, X^{n-1} \rangle_C + \langle X^0, X^1, \dots, X^{n-1} \rangle_C}_{\substack{\subseteq \langle X^0, X^1, \dots, X^{n-1} \rangle_C \\ \text{(since } \langle X^0, X^1, \dots, X^{n-1} \rangle_C \text{ is a } C\text{-module)}}} \\
 &\subseteq X^n + \langle X^0, X^1, \dots, X^{n-1} \rangle_C.
 \end{aligned}$$

In other words, $Q(X) = X^n + W$ for some $W \in \langle X^0, X^1, \dots, X^{n-1} \rangle_C$. Consider this W . We have $W \in \langle X^0, X^1, \dots, X^{n-1} \rangle_C$; thus, we can write W in the form $W = \sum_{i=0}^{n-1} w_i X^i$ for some $w_0, w_1, \dots, w_{n-1} \in C$. Consider these w_0, w_1, \dots, w_{n-1} . Now,

$$Q(X) = X^n + \underbrace{W}_{= \sum_{i=0}^{n-1} w_i X^i} = X^n + \sum_{i=0}^{n-1} w_i X^i.$$

Hence, Q is a monic polynomial with $\deg Q = n$. This proves Lemma 1.8. \square

Proof of Theorem 1.7. Since y is n -integral over A , there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(y) = 0$. Consider this P .

(a) Let C be the A -subalgebra $A[x]$ of B . Then, $C = A[x]$, so that $x \in A[x] = C$.

Now, define a polynomial $Q \in C[X]$ by $Q(X) = P(X - x)$. Then, Lemma 1.8 shows that Q is a monic polynomial with $\deg Q = n$. Also, substituting $x + y$ for

X in the equality $Q(X) = P(X - x)$, we obtain $Q(x + y) = P\left(\underbrace{(x + y) - x}_{=y}\right) = P(y) = 0$.

Hence, there exists a monic polynomial $Q \in C[X]$ with $\deg Q = n$ and $Q(x + y) = 0$. Thus, $x + y$ is n -integral over C . In other words, $x + y$ is n -integral over $A[x]$ (since $C = A[x]$). Thus, Theorem 1.5 (applied to $v = x$ and $u = x + y$) yields that $x + y$ is nm -integral over A . This proves Theorem 1.7 **(a)**.

(b) Recall that $P \in A[X]$ is a monic polynomial with $\deg P = n$. Thus, there exist elements a_0, a_1, \dots, a_{n-1} of A such that $P(X) = X^n + \sum_{k=0}^{n-1} a_k X^k$. Consider

these a_0, a_1, \dots, a_{n-1} . Substituting y for X in $P(X) = X^n + \sum_{k=0}^{n-1} a_k X^k$, we find

$$P(y) = y^n + \sum_{k=0}^{n-1} a_k y^k. \text{ Thus,}$$

$$y^n + \sum_{k=0}^{n-1} a_k y^k = P(y) = 0. \quad (10)$$

Now, define a polynomial $Q \in (A[x])[X]$ by $Q(X) = X^n + \sum_{k=0}^{n-1} x^{n-k} a_k X^k$.

Then,

$$\begin{aligned} Q(xy) &= \underbrace{(xy)^n}_{=x^n y^n} + \sum_{k=0}^{n-1} x^{n-k} \underbrace{a_k (xy)^k}_{\substack{=a_k x^k y^k \\ =x^k a_k y^k}} = x^n y^n + \sum_{k=0}^{n-1} \underbrace{x^{n-k} x^k}_{=x^n} a_k y^k \\ &= x^n y^n + \sum_{k=0}^{n-1} x^n a_k y^k = x^n \underbrace{\left(y^n + \sum_{k=0}^{n-1} a_k y^k \right)}_{\substack{=0 \\ \text{(by (10))}}} = 0. \end{aligned}$$

Also, recall that $Q(X) = X^n + \sum_{k=0}^{n-1} x^{n-k} a_k X^k$; hence, the polynomial $Q \in (A[x])[X]$ is monic and $\deg Q = n$. Thus, there exists a monic polynomial $Q \in (A[x])[X]$ with $\deg Q = n$ and $Q(xy) = 0$. Thus, xy is n -integral over $A[x]$. Hence, Theorem 1.5 (applied to $v = x$ and $u = xy$) yields that xy is nm -integral over A . This proves Theorem 1.7 (b). \square

Corollary 1.9. Let A be a ring. Let B be an A -algebra. Let $x \in B$. Let $m \in \mathbb{N}$. Assume that x is m -integral over A . Then, $-x$ is m -integral over A .

Proof of Corollary 1.9. This is easy to prove directly (using Assertion \mathcal{A} of Theorem 1.1), but the slickest proof is using Theorem 1.7 (b): The element $(-1) \cdot 1_B \in B$ is 1-integral over A (by Theorem 1.6, applied to $a = -1$). Thus, $x \cdot ((-1) \cdot 1_B)$ is $1m$ -integral over A (by Theorem 1.7 (b), applied to $y = (-1) \cdot 1_B$ and $n = 1$). In other words, $-x$ is m -integral over A (since $x \cdot \underbrace{((-1) \cdot 1_B)}_{=-1_B} = x \cdot (-1_B) = -x \cdot 1_B = -x$ and $1m = m$). This proves Corollary 1.9. \square

Corollary 1.10. Let A be a ring. Let B be an A -algebra. Let $x \in B$ and $y \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Assume that x is m -integral over A , and that y is n -integral over A . Then, $x - y$ is nm -integral over A .

Proof of Corollary 1.10. We know that y is n -integral over A . Hence, Corollary 1.9 (applied to y and n instead of x and m) shows that $-y$ is n -integral over A . Thus,

Theorem 1.7 (a) (applied to $-y$ instead of y) shows that $x + (-y)$ is nm -integral over A . In other words, $x - y$ is nm -integral over A (since $x + (-y) = x - y$). This proves Corollary 1.10. \square

1.4. Some further consequences

Theorem 1.11. Let A be a ring. Let B be an A -algebra. Let $n \in \mathbb{N}^+$. Let $v \in B$. Let a_0, a_1, \dots, a_n be $n + 1$ elements of A such that $\sum_{i=0}^n a_i v^i = 0$. Let $k \in \{0, 1, \dots, n\}$. Then, $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -integral over A .

Proof of Theorem 1.11. Let $u = \sum_{i=0}^{n-k} a_{i+k} v^i$. Then,

$$\begin{aligned} 0 &= \sum_{i=0}^n a_i v^i = \sum_{i=0}^{k-1} a_i v^i + \sum_{i=k}^n a_i v^i = \sum_{i=0}^{k-1} a_i v^i + \sum_{i=0}^{n-k} a_{i+k} \underbrace{v^{i+k}}_{=v^i v^k} \\ &\quad \text{(here, we substituted } i+k \text{ for } i \text{ in the second sum)} \\ &= \sum_{i=0}^{k-1} a_i v^i + \underbrace{\sum_{i=0}^{n-k} a_{i+k} v^i v^k}_{=v^k \sum_{i=0}^{n-k} a_{i+k} v^i} = \sum_{i=0}^{k-1} a_i v^i + \underbrace{v^k \sum_{i=0}^{n-k} a_{i+k} v^i}_{=v^k u}, \end{aligned}$$

so that

$$v^k u = - \sum_{i=0}^{k-1} a_i v^i.$$

Let U be the A -submodule $\langle v^0, v^1, \dots, v^{n-1} \rangle_A$ of B . Then, v^0, v^1, \dots, v^{n-1} are n elements of U . Hence, U is an n -generated A -module (since $U = \langle v^0, v^1, \dots, v^{n-1} \rangle_A$). Besides, $n \in \mathbb{N}^+$ and thus $0 \in \{0, 1, \dots, n-1\}$. Therefore, v^0 is one of the n elements v^0, v^1, \dots, v^{n-1} ; hence, $v^0 \in \langle v^0, v^1, \dots, v^{n-1} \rangle_A = U$. Thus, $1 = v^0 \in U$.

Note that U is an A -submodule of B , and thus is closed under A -linear combination.

Now, we are going to show that

$$uv^s \in U \quad \text{for any } s \in \{0, 1, \dots, n-1\}. \quad (11)$$

[*Proof of (11).* Let $s \in \{0, 1, \dots, n-1\}$. Thus, we are in one of the following two cases:

Case 1: We have $s < k$.

Case 2: We have $s \geq k$.

Let us first consider Case 1. In this case, we have $s < k$. Hence, $s \leq k-1$ (since s and k are integers) and thus $s \in \{0, 1, \dots, k-1\}$ (since $s \in \{0, 1, \dots, n-1\}$).

For each $i \in \{0, 1, \dots, n-k\}$, we have $i \leq n-k$ and thus $\underbrace{i}_{\leq n-k} + \underbrace{s}_{\leq k-1} \leq (n-k) + (k-1) = n-1$ and therefore $i+s \in \{0, 1, \dots, n-1\}$ (since $\underbrace{i}_{\geq 0} + \underbrace{s}_{\geq 0} \geq 0$) and therefore

$$v^{i+s} \in \{v^0, v^1, \dots, v^{n-1}\} \subseteq \langle v^0, v^1, \dots, v^{n-1} \rangle_A = U.$$

Hence, $\sum_{i=0}^{n-k} a_{i+k} v^{i+s}$ is an A -linear combination of elements of the set U (since the coefficients a_{i+k} belong to A) and therefore belongs to U itself (since U is closed under A -linear combination). In other words, $\sum_{i=0}^{n-k} a_{i+k} v^{i+s} \in U$.

Now, from $u = \sum_{i=0}^{n-k} a_{i+k} v^i$, we obtain

$$uv^s = \sum_{i=0}^{n-k} a_{i+k} \underbrace{v^i \cdot v^s}_{=v^{i+s}} = \sum_{i=0}^{n-k} a_{i+k} v^{i+s} \in U.$$

Hence, (11) is proven in Case 1.

Let us next consider Case 2. In this case, we have $s \geq k$. Hence, $s-k \geq 0$. Also, $s \leq n-1$ (since $s \in \{0, 1, \dots, n-1\}$).

For each $i \in \{0, 1, \dots, k-1\}$, we have $i \geq 0$ and $i \leq k-1$ and thus $\underbrace{i}_{\leq k-1} + (s-k) \leq (k-1) + (s-k) = s-1 \leq s \leq n-1$ and therefore $i+(s-k) \in \{0, 1, \dots, n-1\}$ (since $\underbrace{i}_{\geq 0} + (s-k) \geq s-k \geq 0$) and thus

$$v^{i+(s-k)} \in \{v^0, v^1, \dots, v^{n-1}\} \subseteq \langle v^0, v^1, \dots, v^{n-1} \rangle_A = U.$$

Hence, $\sum_{i=0}^{k-1} a_i v^{i+(s-k)}$ is an A -linear combination of elements of the set U (since the coefficients a_i belong to A) and therefore belongs to U itself (since U is closed under A -linear combination). In other words, $\sum_{i=0}^{k-1} a_i v^{i+(s-k)} \in U$.

From $s-k \geq 0$, we obtain $s-k \in \mathbb{N}$ and thus $v^s = v^{k+(s-k)} = v^k v^{s-k}$. Hence,

$$\begin{aligned} uv^s &= uv^k v^{s-k} = v^k u \cdot v^{s-k} = - \sum_{i=0}^{k-1} a_i \underbrace{v^i \cdot v^{s-k}}_{=v^{i+(s-k)}} \quad \left(\text{since } v^k u = - \sum_{i=0}^{k-1} a_i v^i \right) \\ &= - \underbrace{\sum_{i=0}^{k-1} a_i v^{i+(s-k)}}_{\in U} \in -U \subseteq U \quad (\text{since } U \text{ is an } A\text{-module}). \end{aligned}$$

Hence, (11) is proven in Case 2.

Hence, in both cases, we have proven (11). This completes the proof of (11).]

Thus we know that $uv^s \in U$ for every $s \in \{0, 1, \dots, n-1\}$. Hence, Lemma 0.3 (applied to $M = B$, $N = U$, $S = \{0, 1, \dots, n-1\}$ and $m_s = uv^s$) yields

$$\langle uv^s \mid s \in \{0, 1, \dots, n-1\} \rangle_A \subseteq U.$$

Now, from $U = \langle v^0, v^1, \dots, v^{n-1} \rangle_A$, we obtain

$$\begin{aligned} uU &= u \langle v^0, v^1, \dots, v^{n-1} \rangle_A = \langle uv^0, uv^1, \dots, uv^{n-1} \rangle_A \\ &= \langle uv^s \mid s \in \{0, 1, \dots, n-1\} \rangle_A \subseteq U. \end{aligned}$$

Altogether, U is an n -generated A -submodule of B such that $1 \in U$ and $uU \subseteq U$. Thus, $u \in B$ satisfies Assertion \mathcal{C} of Theorem 1.1. Hence, $u \in B$ satisfies the four equivalent assertions \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} of Theorem 1.1. Consequently, u is n -integral over A . Since $u = \sum_{i=0}^{n-k} a_{i+k} v^i$, this means that $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -integral over A . This proves Theorem 1.11. \square

Corollary 1.12. Let A be a ring. Let B be an A -algebra. Let $\alpha \in \mathbb{N}$ and $\beta \in \mathbb{N}$ be such that $\alpha + \beta \in \mathbb{N}^+$. Let $u \in B$ and $v \in B$. Let $s_0, s_1, \dots, s_\alpha$ be $\alpha + 1$ elements of A such that $\sum_{i=0}^{\alpha} s_i v^i = u$. Let t_0, t_1, \dots, t_β be $\beta + 1$ elements of A such that $\sum_{i=0}^{\beta} t_i v^{\beta-i} = uv^\beta$. Then, u is $(\alpha + \beta)$ -integral over A .

(This Corollary 1.12 generalizes [4, Exercise 2-5], which says that if v is an invertible element of an A -algebra B , then every element $u \in A[v] \cap A[v^{-1}]$ is integral over A . To see how this follows from Corollary 1.12, just pick $\alpha \in \mathbb{N}^+$ and $\beta \in \mathbb{N}^+$ and $s_0, s_1, \dots, s_\alpha \in A$ and $t_0, t_1, \dots, t_\beta \in A$ such that $\sum_{i=0}^{\alpha} s_i v^i = u$ and

$$\sum_{i=0}^{\beta} t_i (v^{-1})^i = u.)$$

First proof of Corollary 1.12. Let $k = \beta$ and $n = \alpha + \beta$. Then, $k \in \{0, 1, \dots, n\}$ (since $\alpha \in \mathbb{N}$ and $\beta \in \mathbb{N}$) and $n = \alpha + \beta \in \mathbb{N}^+$ and $n - \beta = \alpha$ (since $n = \alpha + \beta$). Define $n + 1$ elements a_0, a_1, \dots, a_n of A by

$$a_i = \begin{cases} t_{\beta-i}, & \text{if } i < \beta; \\ t_0 - s_0, & \text{if } i = \beta; \\ -s_{i-\beta}, & \text{if } i > \beta \end{cases} \quad \text{for every } i \in \{0, 1, \dots, n\}.$$

Then, from $n = \alpha + \beta$, we obtain

$$\begin{aligned}
 \sum_{i=0}^n a_i v^i &= \sum_{i=0}^{\alpha+\beta} a_i v^i = \sum_{i=0}^{\beta-1} \underbrace{a_i}_{=t_{\beta-i}} v^i + \sum_{i=\beta}^{\beta} \underbrace{a_i}_{=t_0-s_0} v^i + \sum_{i=\beta+1}^{\alpha+\beta} \underbrace{a_i}_{=-s_{i-\beta}} v^i \\
 &\quad \text{(by the definition of } a_i, \text{ since } i < \beta) \quad \text{(by the definition of } a_i, \text{ since } i = \beta) \quad \text{(by the definition of } a_i, \text{ since } i > \beta) \\
 &= \sum_{i=0}^{\beta-1} t_{\beta-i} v^i + \underbrace{\sum_{i=\beta}^{\beta} (t_0 - s_0) v^i}_{=(t_0-s_0)v^\beta} + \underbrace{\sum_{i=\beta+1}^{\alpha+\beta} (-s_{i-\beta}) v^i}_{=-\sum_{i=\beta+1}^{\alpha+\beta} s_{i-\beta} v^i} \\
 &= \sum_{i=0}^{\beta-1} t_{\beta-i} v^i + t_0 v^\beta - s_0 v^\beta - \sum_{i=\beta+1}^{\alpha+\beta} s_{i-\beta} v^i \\
 &= \sum_{i=0}^{\beta-1} t_{\beta-i} v^i + t_0 v^\beta - \left(s_0 v^\beta + \sum_{i=\beta+1}^{\alpha+\beta} s_{i-\beta} v^i \right) \\
 &= \sum_{i=0}^{\beta-1} t_{\beta-i} v^i + t_0 v^\beta - \left(s_0 v^\beta + \sum_{i=1}^{\alpha} \underbrace{s_{(i+\beta)-\beta}}_{=s_i} \underbrace{v^{i+\beta}}_{=v^i v^\beta} \right) \\
 &\quad \text{(here, we substituted } i + \beta \text{ for } i \text{ in the second sum)} \\
 &= \sum_{i=0}^{\beta-1} t_{\beta-i} v^i + t_0 v^\beta - \left(s_0 v^\beta + \sum_{i=1}^{\alpha} s_i v^i v^\beta \right) \\
 &= \sum_{i=1}^{\beta} \underbrace{t_{\beta-(\beta-i)}}_{=t_i} v^{\beta-i} + t_0 \underbrace{v^\beta}_{=v^{\beta-0}} - \left(s_0 \underbrace{v^\beta}_{=v^0 v^\beta} + \sum_{i=1}^{\alpha} s_i v^i v^\beta \right) \\
 &\quad \text{(here, we substituted } \beta - i \text{ for } i \text{ in the first sum)} \\
 &= \sum_{i=1}^{\beta} t_i v^{\beta-i} + t_0 v^{\beta-0} - \left(s_0 v^0 v^\beta + \sum_{i=1}^{\alpha} s_i v^i v^\beta \right) \\
 &= \underbrace{\sum_{i=1}^{\beta} t_i v^{\beta-i} + t_0 v^{\beta-0}}_{=\sum_{i=0}^{\beta} t_i v^{\beta-i} = u v^\beta} - \left(\underbrace{s_0 v^0 + \sum_{i=1}^{\alpha} s_i v^i}_{=\sum_{i=0}^{\alpha} s_i v^i = u} \right) v^\beta = u v^\beta - u v^\beta = 0.
 \end{aligned}$$

Thus, Theorem 1.11 yields that $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -integral over A .

But $k = \beta$ and thus

$$\begin{aligned}
 \sum_{i=0}^{n-k} a_{i+k} v^i &= \sum_{i=0}^{n-\beta} a_{i+\beta} v^i = \sum_{i=\beta}^n \underbrace{a_{(i-\beta)+\beta}}_{=a_i} v^{i-\beta} \\
 &\quad \text{(here, we have substituted } i - \beta \text{ for } i \text{ in the sum)} \\
 &= \sum_{i=\beta}^n a_i v^{i-\beta} = \sum_{i=\beta}^{\beta} \underbrace{a_i}_{\substack{=t_0-s_0 \\ \text{(by the} \\ \text{definition of } a_i, \\ \text{since } i=\beta)}} v^{i-\beta} + \sum_{i=\beta+1}^n \underbrace{a_i}_{\substack{=-s_{i-\beta} \\ \text{(by the} \\ \text{definition of } a_i, \\ \text{since } i>\beta)}} v^{i-\beta} \\
 &= \underbrace{\sum_{i=\beta}^{\beta} (t_0 - s_0) v^{i-\beta}}_{=(t_0-s_0)v^{\beta-\beta}} + \underbrace{\sum_{i=\beta+1}^n (-s_{i-\beta}) v^{i-\beta}}_{=\sum_{i=1}^{n-\beta} (-s_i) v^i} \\
 &\quad \text{(here, we have substituted } i \\
 &\quad \text{for } i-\beta \text{ in the sum)} \\
 &= (t_0 - s_0) \underbrace{v^{\beta-\beta}}_{=v^0} + \underbrace{\sum_{i=1}^{n-\beta} (-s_i) v^i}_{=-\sum_{i=1}^{n-\beta} s_i v^i} = \underbrace{(t_0 - s_0) v^0}_{=t_0 v^0 - s_0 v^0} + \left(-\sum_{i=1}^{\alpha} s_i v^i \right) \\
 &\quad \text{(since } n-\beta=\alpha) \\
 &= t_0 v^0 - s_0 v^0 + \left(-\sum_{i=1}^{\alpha} s_i v^i \right) = t_0 \underbrace{v^0}_{=1_B} - \underbrace{\left(s_0 v^0 + \sum_{i=1}^{\alpha} s_i v^i \right)}_{=\sum_{i=0}^{\alpha} s_i v^i = u} \\
 &= t_0 \cdot 1_B - u.
 \end{aligned}$$

Thus, $t_0 \cdot 1_B - u$ is n -integral over A (since $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -integral over A).

Thus, Corollary 1.9 (applied to $x = t_0 \cdot 1_B - u$ and $m = n$) shows that $-(t_0 \cdot 1_B - u)$ is n -integral over A . In other words, $u - t_0 \cdot 1_B$ is n -integral over A (since $-(t_0 \cdot 1_B - u) = u - t_0 \cdot 1_B$).

On the other hand, $t_0 \cdot 1_B$ is 1-integral over A (by Theorem 1.6, applied to $a = t_0$). Thus, $t_0 \cdot 1_B + (u - t_0 \cdot 1_B)$ is $n \cdot 1$ -integral over A (by Theorem 1.7 (a), applied to $x = t_0 \cdot 1_B$, $y = u - t_0 \cdot 1_B$ and $m = 1$). In other words, u is $(\alpha + \beta)$ -integral over A (since $t_0 \cdot 1_B + (u - t_0 \cdot 1_B) = u$ and $n \cdot 1 = n = \alpha + \beta$). This proves Corollary 1.12. \square

We will provide a second proof of Corollary 1.12 in Section 5.

Corollary 1.13. Let A be a ring. Let B be an A -algebra. Let $n \in \mathbb{N}^+$ and $m \in \mathbb{N}$. Let $v \in B$. Let b_0, b_1, \dots, b_{n-1} be n elements of A , and let $u = \sum_{i=0}^{n-1} b_i v^i$. Assume that vu is m -integral over A . Then, u is nm -integral over A .

Corollary 1.13 generalizes a folklore fact about integrality, which states that if B is an A -algebra, and if an invertible $v \in B$ satisfies $v^{-1} \in A[v]$, then v is integral over A . (Indeed, this latter fact follows from Corollary 1.13 by setting $u = v^{-1}$.)

Proof of Corollary 1.13. Define $n + 1$ elements a_0, a_1, \dots, a_n of $A[vu]$ by

$$a_i = \begin{cases} -vu, & \text{if } i = 0; \\ b_{i-1} \cdot 1_B, & \text{if } i > 0 \end{cases} \quad \text{for every } i \in \{0, 1, \dots, n\}.$$

(These are well-defined, since every positive $i \in \{0, 1, \dots, n\}$ satisfies $i \in \{1, 2, \dots, n\}$ and thus $i - 1 \in \{0, 1, \dots, n - 1\}$ and thus $b_{i-1} \in A$ and therefore $b_{i-1} \cdot 1_B \in A \cdot 1_B \subseteq A[vu]$.)

The definition of a_0 yields $a_0 = -vu$. Also,

$$\begin{aligned} \sum_{i=0}^n a_i v^i &= \underbrace{a_0}_{=-vu} \underbrace{v^0}_{=1} + \sum_{i=1}^n \underbrace{a_i}_{=b_{i-1} \cdot 1_B} v^i = -vu + \sum_{i=1}^n b_{i-1} \cdot \underbrace{1_B v^i}_{=v^i = v^{i-1} v} \\ &= -vu + \sum_{i=1}^n b_{i-1} v^{i-1} v = -vu + \underbrace{\sum_{i=0}^{n-1} b_i v^i}_{=u} \\ &\quad \text{(here, we substituted } i \text{ for } i - 1 \text{ in the sum)} \\ &= -vu + uv = 0. \end{aligned}$$

Let $k = 1$. Then, $k = 1 \in \{0, 1, \dots, n\}$ (since $n \in \mathbb{N}^+$).

Now, $A[vu]$ is a subring of B ; hence, B is an $A[vu]$ -algebra. The $n + 1$ elements a_0, a_1, \dots, a_n of $A[vu]$ satisfy $\sum_{i=0}^n a_i v^i = 0$.

Hence, Theorem 1.11 (applied to the ring $A[vu]$ in lieu of A) yields that $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -integral over $A[vu]$. But from $k = 1$, we obtain

$$\sum_{i=0}^{n-k} a_{i+k} v^i = \sum_{i=0}^{n-1} \underbrace{a_{i+1}}_{=b_{(i+1)-1} \cdot 1_B} v^i = \sum_{i=0}^{n-1} \underbrace{b_{(i+1)-1}}_{=b_i} \cdot \underbrace{1_B v^i}_{=v^i} = \sum_{i=0}^{n-1} b_i v^i = u.$$

(by the definition of a_{i+1} , since $i+1 > 0$)

Hence, u is n -integral over $A[vu]$ (since $\sum_{i=0}^{n-k} a_{i+k}v^i$ is n -integral over $A[vu]$). But vu is m -integral over A . Thus, Theorem 1.5 (applied to vu in lieu of v) yields that u is nm -integral over A . This proves Corollary 1.13. \square

2. Integrality over ideal semifiltrations

2.1. Definitions of ideal semifiltrations and integrality over them

We now set our sights at a more general notion of integrality.

Definition 2.1. Let A be a ring, and let $(I_\rho)_{\rho \in \mathbb{N}}$ be a sequence of ideals of A . Then, $(I_\rho)_{\rho \in \mathbb{N}}$ is called an *ideal semifiltration* of A if and only if it satisfies the two conditions

$$\begin{aligned} I_0 &= A; \\ I_a I_b &\subseteq I_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

Two simple examples of ideal semifiltrations can easily be constructed from any ideal:

Example 2.2. Let A be a ring. Let I be an ideal of A . Then:

(a) The sequence $(I^\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A . (Here, I^ρ denotes the ρ -th power of I in the multiplicative monoid of ideals of A ; in particular, $I^0 = A$.)

(b) The sequence $(A, I, I, I, \dots) = \left(\begin{cases} A, & \text{if } \rho = 0; \\ I, & \text{if } \rho > 0 \end{cases} \right)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A .

Proof of Example 2.2. This is a straightforward exercise in checking axioms. \square

Definition 2.3. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $n \in \mathbb{N}$. Let $u \in B$.

We say that the element u of B is *n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$* if there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

This definition generalizes [5, Definition 1.1.1] in multiple ways. Indeed, if I is an ideal of a ring A , and if $u \in A$ and $n \in \mathbb{N}$, then u is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$ (here, $(I_\rho)_{\rho \in \mathbb{N}}$ is the ideal semifiltration from Example 2.2 (a)) if and only if there is an equation of integral dependence of u over I (in the sense of [5, Definition 1.1.1]).

We further notice that integrality over an ideal semifiltration of a ring A is a stronger claim than integrality over A :

Proposition 2.4. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $n \in \mathbb{N}$. Let $u \in B$ be such that u is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. Then, u is n -integral over A .

Proof of Proposition 2.4. We know that u is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. Thus, by Definition 2.3, there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (a_0, a_1, \dots, a_n) .

For each $k \in \{0, 1, \dots, n\}$, we have $a_k \in I_{n-k}$ (since $a_i \in I_{n-i}$ for every $i \in \{0, 1, \dots, n\}$) and therefore $a_k \in I_{n-k} \subseteq A$. Thus, we can define a polynomial $P \in A[X]$ by $P(X) = \sum_{k=0}^n a_k X^k$. Consider this P . This polynomial P satisfies $\deg P \leq n$, and its coefficient before X^n is $a_n = 1$. Hence, this polynomial P is monic and satisfies $\deg P = n$. Also, by substituting u for X in $P(X) = \sum_{k=0}^n a_k X^k$,

we obtain $P(u) = \sum_{k=0}^n a_k u^k = 0$. Hence, we have found a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$.

In other words, u is n -integral over A . This proves Proposition 2.4. \square

We leave it to the reader to prove the following simple fact, which shows that nilpotency is an instance of integrality over ideal semifiltrations:

Proposition 2.5. Let A be a ring. Let $0A$ be the zero ideal of A . Let $n \in \mathbb{N}$. Let $u \in A$. Then, the element u of A is n -integral over $(A, ((0A)^\rho)_{\rho \in \mathbb{N}})$ if and only if $u^n = 0$.

2.2. Polynomial rings and Rees algebras

In order to study integrality over ideal semifiltrations, we shall now introduce the concept of a Rees algebra – a subalgebra of a polynomial ring that conveniently encodes an ideal semifiltration of the base ring. This, again, generalizes

a classical notion for ideals (namely, the Rees algebra of an ideal – see [5, Definition 5.1.1]). First, we recall a basic fact:

Definition 2.6. Let A be a ring. Let B be an A -algebra. Then, there is a canonical ring homomorphism $\iota : A \rightarrow B$ that sends each $a \in A$ to $a \cdot 1_B \in B$. This ring homomorphism ι induces a canonical ring homomorphism $\iota[Y] : A[Y] \rightarrow B[Y]$ between the polynomial rings $A[Y]$ and $B[Y]$ that sends each polynomial $\sum_{i=0}^m a_i Y^i \in A[Y]$ (with $m \in \mathbb{N}$ and $(a_0, a_1, \dots, a_m) \in A^{m+1}$) to the polynomial $\sum_{i=0}^m \iota(a_i) Y^i \in B[Y]$. Thus, the polynomial ring $B[Y]$ becomes an $A[Y]$ -algebra.

Definition 2.7. Let A be a ring, and let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Thus, I_0, I_1, I_2, \dots are ideals of A , and we have $I_0 = A$.

Consider the polynomial ring $A[Y]$. For each $i \in \mathbb{N}$, the subset $I_i Y^i$ of $A[Y]$ is an A -submodule of the A -algebra $A[Y]$ (since I_i is an ideal of A). Hence, the sum $\sum_{i \in \mathbb{N}} I_i Y^i$ of these A -submodules must also be an A -submodule of the A -algebra $A[Y]$.

Let $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ denote this A -submodule $\sum_{i \in \mathbb{N}} I_i Y^i$ of the A -algebra $A[Y]$. Then,

$$\begin{aligned} A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] &= \sum_{i \in \mathbb{N}} I_i Y^i \\ &= \left\{ \sum_{i \in \mathbb{N}} a_i Y^i \mid (a_i \in I_i \text{ for all } i \in \mathbb{N}), \right. \\ &\quad \left. \text{and (only finitely many } i \in \mathbb{N} \text{ satisfy } a_i \neq 0) \right\} \\ &= \{ P \in A[Y] \mid \text{the } i\text{-th coefficient of the polynomial } P \\ &\quad \text{lies in } I_i \text{ for every } i \in \mathbb{N} \}. \end{aligned}$$

Clearly, $A \subseteq A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$, since

$$A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] = \sum_{i \in \mathbb{N}} I_i Y^i \supseteq \underbrace{I_0}_{=A} \underbrace{Y^0}_{=1} = A \cdot 1 = A.$$

Hence, $1 \in A \subseteq A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Also, the A -submodule $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ of $A[Y]$ is an A -subalgebra of the A -algebra $A[Y]$ (by Lemma 2.8 below), and thus is a subring of $A[Y]$.

This A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is called the *Rees algebra* of the ideal semifiltration $(I_\rho)_{\rho \in \mathbb{N}}$.

Lemma 2.8. Let A be a ring, and let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Then, the A -submodule $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ of $A[Y]$ is an A -subalgebra of the A -algebra $A[Y]$.

Proof of Lemma 2.8. Multiplying the equality $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] = \sum_{i \in \mathbb{N}} I_i Y^i$ with itself, we find

$$\begin{aligned}
 & A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \cdot A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \\
 &= \left(\sum_{i \in \mathbb{N}} I_i Y^i \right) \cdot \left(\sum_{i \in \mathbb{N}} I_i Y^i \right) = \left(\sum_{i \in \mathbb{N}} I_i Y^i \right) \cdot \left(\sum_{j \in \mathbb{N}} I_j Y^j \right) \\
 &\quad \text{(here we renamed the index } i \text{ as } j \text{ in the second sum)} \\
 &= \sum_{i \in \mathbb{N}} \sum_{j \in \mathbb{N}} \underbrace{I_i Y^i I_j Y^j}_{= I_j Y^i} = \sum_{i \in \mathbb{N}} \sum_{j \in \mathbb{N}} \underbrace{I_i I_j}_{\subseteq I_{i+j}} \underbrace{Y^i Y^j}_{= Y^{i+j}} \\
 &\quad \text{(since } (I_\rho)_{\rho \in \mathbb{N}} \text{ is an ideal semifiltration)} \\
 &\subseteq \sum_{i \in \mathbb{N}} \sum_{j \in \mathbb{N}} \underbrace{I_{i+j} Y^{i+j}}_{\subseteq \sum_{k \in \mathbb{N}} I_k Y^k} \subseteq \sum_{i \in \mathbb{N}} \sum_{j \in \mathbb{N}} \sum_{k \in \mathbb{N}} I_k Y^k \\
 &\subseteq \sum_{k \in \mathbb{N}} I_k Y^k \quad \left(\text{since } \sum_{k \in \mathbb{N}} I_k Y^k \text{ is an } A\text{-module} \right) \\
 &= \sum_{i \in \mathbb{N}} I_i Y^i \quad \text{(here we renamed the index } k \text{ as } i \text{ in the sum)} \\
 &= A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right].
 \end{aligned}$$

Hence, the A -submodule $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ of $A[Y]$ is closed under multiplication. Thus, it is an A -subalgebra of the A -algebra $A[Y]$ (since $1 \in A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$). This proves Lemma 2.8. \square

Remark 2.9. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A .

Consider the polynomial ring $A[Y]$ and its A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ defined in Definition 2.7.

The polynomial ring $B[Y]$ is an $A[Y]$ -algebra (as explained in Definition 2.6), and thus is an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra (since $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is a subring of $A[Y]$). Hence, if $p \in B[Y]$ is a polynomial and $n \in \mathbb{N}$, then it makes sense to ask whether p is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Questions of this form will often appear in what follows.

We note in passing that the notion of a Rees algebra helps set up a bijection between ideal semifiltrations of a ring A and graded A -subalgebras of the polynomial ring $A[Y]$:

Proposition 2.10. Let A be a ring. Consider the polynomial ring $A[Y]$ as a graded A -algebra (with the usual degree of polynomials).

(a) If $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , then its Rees algebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is a graded A -subalgebra of $A[Y]$.

(b) If B is any graded A -subalgebra of $A[Y]$, and if $\rho \in \mathbb{N}$, then we let $I_{B,\rho}$ denote the subset $\{a \in A \mid aY^\rho \in B\}$ of A . Then, $I_{B,\rho}$ is an ideal of A .

(c) The maps

$$\begin{aligned} \{\text{ideal semifiltrations of } A\} &\rightarrow \{\text{graded } A\text{-subalgebras of } A[Y]\}, \\ (I_\rho)_{\rho \in \mathbb{N}} &\mapsto A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \end{aligned}$$

and

$$\begin{aligned} \{\text{graded } A\text{-subalgebras of } A[Y]\} &\rightarrow \{\text{ideal semifiltrations of } A\}, \\ B &\mapsto (I_{B,\rho})_{\rho \in \mathbb{N}} \end{aligned}$$

are mutually inverse bijections.

We shall not have any need for this proposition, so we omit its (straightforward and easy) proof.

2.3. Reduction to integrality over rings

We start with a theorem which reduces the question of n -integrality over $(A, (I_\rho)_{\rho \in \mathbb{N}})$ to that of n -integrality over a ring³:

Theorem 2.11. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $n \in \mathbb{N}$. Let $u \in B$.

Consider the polynomial ring $A[Y]$ and its A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ defined in Definition 2.7.

Then, the element u of B is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$ if and only if the element uY of the polynomial ring $B[Y]$ is n -integral over the ring $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. (Here, $B[Y]$ is an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra, as explained in Remark 2.9.)

Proof of Theorem 2.11. In order to verify Theorem 2.11, we have to prove the following two lemmata:

³Theorem 2.11 is inspired by [5, Proposition 5.2.1].

Lemma \mathcal{E} : If u is n -integral over $\left(A, (I_\rho)_{\rho \in \mathbb{N}}\right)$, then uY is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$.

Lemma \mathcal{F} : If uY is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$, then u is n -integral over $\left(A, (I_\rho)_{\rho \in \mathbb{N}}\right)$.

[*Proof of Lemma \mathcal{E} :* Assume that u is n -integral over $\left(A, (I_\rho)_{\rho \in \mathbb{N}}\right)$. Thus, by Definition 2.3, there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (a_0, a_1, \dots, a_n) .

For each $k \in \{0, 1, \dots, n\}$, we have $a_k \in I_{n-k}$ (since $a_i \in I_{n-i}$ for every $i \in \{0, 1, \dots, n\}$) and therefore

$$\underbrace{a_k}_{\in I_{n-k}} Y^{n-k} \in I_{n-k} Y^{n-k} \subseteq \sum_{i \in \mathbb{N}} I_i Y^i = A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$$

(since Definition 2.7 yields $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] = \sum_{i \in \mathbb{N}} I_i Y^i$).

Thus, we can define a polynomial $P \in \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right) [X]$ by $P(X) = \sum_{k=0}^n a_k Y^{n-k} X^k$. Consider this P . This polynomial P satisfies $\deg P \leq n$, and its coefficient before X^n is $\underbrace{a_n}_{=1} \underbrace{Y^{n-n}}_{=Y^0=1} = 1$. Hence, this polynomial P is monic and

satisfies $\deg P = n$. Also, by substituting uY for X in $P(X) = \sum_{k=0}^n a_k Y^{n-k} X^k$, we obtain

$$P(uY) = \sum_{k=0}^n a_k Y^{n-k} \underbrace{(uY)^k}_{=u^k Y^k} = \sum_{k=0}^n a_k Y^{n-k} u^k Y^k = \sum_{k=0}^n a_k u^k \underbrace{Y^{n-k} Y^k}_{=Y^n} = Y^n \cdot \underbrace{\sum_{k=0}^n a_k u^k}_{=0} = 0.$$

Thus, there exists a monic polynomial $P \in \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right) [X]$ with $\deg P = n$ and $P(uY) = 0$. Hence, uY is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. This proves Lemma \mathcal{E} .]

[*Proof of Lemma \mathcal{F} :* Assume that uY is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Thus, there exists a monic polynomial $P \in \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right) [X]$ with $\deg P = n$

and $P(uY) = 0$. Consider this P . Since $P \in \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right) [X]$ satisfies $\deg P = n$, there exists $(p_0, p_1, \dots, p_n) \in \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right)^{n+1}$ such that $P(X) = \sum_{k=0}^n p_k X^k$. Consider this (p_0, p_1, \dots, p_n) . Note that $p_n = 1$ (since P is monic and $\deg P = n$).

Recall that $(p_0, p_1, \dots, p_n) \in \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right)^{n+1}$. Hence, for every $k \in \{0, 1, \dots, n\}$, we have $p_k \in A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] = \sum_{i \in \mathbb{N}} I_i Y^i$, and thus there exists a sequence $(p_{k,i})_{i \in \mathbb{N}} \in A^{\mathbb{N}}$ such that $p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i$, such that $(p_{k,i} \in I_i \text{ for every } i \in \mathbb{N})$, and such that only finitely many $i \in \mathbb{N}$ satisfy $p_{k,i} \neq 0$. Consider this sequence. Thus,

$$P(X) = \sum_{k=0}^n \underbrace{p_k}_{=\sum_{i \in \mathbb{N}} p_{k,i} Y^i} X^k = \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} Y^i X^k.$$

Substituting uY for X in this equality, we find

$$\begin{aligned} P(uY) &= \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} Y^i \underbrace{(uY)^k}_{\substack{=u^k Y^k \\ =Y^k u^k}} = \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} \underbrace{Y^i Y^k}_{=Y^{i+k}} u^k \\ &= \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} Y^{i+k} u^k = \sum_{k \in \{0,1,\dots,n\}} \sum_{i \in \mathbb{N}} p_{k,i} Y^{i+k} u^k \\ &= \sum_{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}} p_{k,i} Y^{i+k} u^k = \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} \underbrace{Y^{i+k}}_{=Y^\ell} u^k \\ &= \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} Y^\ell u^k = \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} u^k Y^\ell. \end{aligned}$$

Comparing this with $P(uY) = 0$, we find $\sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} u^k Y^\ell = 0$.

In other words, the polynomial $\sum_{\ell \in \mathbb{N}} \underbrace{\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} u^k Y^\ell}_{\in B} \in B[Y]$ equals 0.

Hence, its coefficient before Y^n equals 0 as well. But its coefficient before Y^n is $\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=n}} p_{k,i} u^k$. Comparing the preceding two sentences, we see that

$\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=n}} p_{k,i} u^k$ equals 0. Thus,

$$0 = \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=n}} p_{k,i} u^k = \sum_{k \in \{0,1,\dots,n\}} \sum_{\substack{i \in \mathbb{N}; \\ i+k=n}} p_{k,i} u^k. \quad (12)$$

For each $k \in \{0,1,\dots,n\}$, we have

$$\left\{ i \in \mathbb{N} \mid \underbrace{i+k=n}_{\Leftrightarrow (i=n-k)} \right\} = \{i \in \mathbb{N} \mid i = n - k\} = \{n - k\}$$

(because $n - k \in \mathbb{N}$ (since $k \in \{0,1,\dots,n\}$)) and thus

$$\sum_{\substack{i \in \mathbb{N}; \\ i+k=n}} p_{k,i} u^k = \sum_{i \in \{n-k\}} p_{k,i} u^k = p_{k,n-k} u^k.$$

Thus, (12) becomes

$$0 = \sum_{k \in \{0,1,\dots,n\}} \underbrace{\sum_{\substack{i \in \mathbb{N}; \\ i+k=n}} p_{k,i} u^k}_{=p_{k,n-k} u^k} = \sum_{k \in \{0,1,\dots,n\}} p_{k,n-k} u^k. \quad (13)$$

Recall that $p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i$ for every $k \in \{0,1,\dots,n\}$. Applying this to $k = n$, we find $p_n = \sum_{i \in \mathbb{N}} p_{n,i} Y^i$. Comparing this with $p_n = 1 = 1 \cdot Y^0$, we find

$$\sum_{i \in \mathbb{N}} p_{n,i} Y^i = 1 \cdot Y^0 \quad \text{in } A[Y].$$

Hence, the coefficient of the polynomial $\sum_{i \in \mathbb{N}} p_{n,i} Y^i \in A[Y]$ before Y^0 is 1. But the coefficient of the polynomial $\sum_{i \in \mathbb{N}} p_{n,i} Y^i \in A[Y]$ before Y^0 is $p_{n,0}$ (since $p_{n,i} \in A$ for all $i \in \mathbb{N}$). Comparing the preceding two sentences, we see that $p_{n,0} = 1$.

Define an $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ by setting

$$(a_k = p_{k,n-k} \text{ for every } k \in \{0,1,\dots,n\}).$$

Then, $a_n = p_{n,n-n} = p_{n,0} = 1$. Besides,

$$\sum_{k=0}^n \underbrace{a_k}_{\substack{=p_{k,n-k} \\ \text{(by the definition} \\ \text{of } a_k)}} u^k = \sum_{k=0}^n p_{k,n-k} u^k = \sum_{k \in \{0,1,\dots,n\}} p_{k,n-k} u^k = 0$$

(by (13)). Finally, for every $k \in \{0, 1, \dots, n\}$, we have $n - k \in \mathbb{N}$ and thus $a_k = p_{k, n-k} \in I_{n-k}$ (since $p_{k,i} \in I_i$ for every $i \in \mathbb{N}$). Renaming the variable k as i in this statement, we obtain the following: For every $i \in \{0, 1, \dots, n\}$, we have $a_i \in I_{n-i}$.

Altogether, we now know that the $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ satisfies

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Thus, by Definition 2.3, the element u is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. This proves Lemma \mathcal{F} .]

Combining Lemma \mathcal{E} and Lemma \mathcal{F} , we obtain that u is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$ if and only if uY is n -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. This proves Theorem 2.11. \square

2.4. Sums and products again

Let us next state an analogue of Theorem 1.6 for integrality over ideal semifiltrations:

Theorem 2.12. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $u \in A$. Then, $u \cdot 1_B$ is 1-integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$ if and only if $u \cdot 1_B \in I_1 \cdot 1_B$.

Proof of Theorem 2.12. In order to verify Theorem 2.12, we have to prove the following two lemmata:

Lemma \mathcal{G} : If $u \cdot 1_B$ is 1-integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$, then $u \cdot 1_B \in I_1 \cdot 1_B$.

Lemma \mathcal{H} : If $u \cdot 1_B \in I_1 \cdot 1_B$, then $u \cdot 1_B$ is 1-integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$.

[*Proof of Lemma \mathcal{G} :* Assume that $u \cdot 1_B$ is 1-integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $u \cdot 1_B$ and 1 instead of u and n), there exists some $(a_0, a_1) \in A^2$ such that

$$\sum_{k=0}^1 a_k (u \cdot 1_B)^k = 0, \quad a_1 = 1, \quad \text{and} \quad a_i \in I_{1-i} \text{ for every } i \in \{0, 1\}.$$

Consider this (a_0, a_1) . Thus, $a_0 \in I_{1-0}$ (since $a_i \in I_{1-i}$ for every $i \in \{0, 1\}$), so that $a_0 \in I_{1-0} = I_1$ and thus $-a_0 \in -I_1 \subseteq I_1$ (since I_1 is an ideal of A). Also,

$$0 = \sum_{k=0}^1 a_k (u \cdot 1_B)^k = a_0 \underbrace{(u \cdot 1_B)^0}_{=1_B} + \underbrace{a_1}_{=1} \underbrace{(u \cdot 1_B)^1}_{=u \cdot 1_B} = a_0 \cdot 1_B + u \cdot 1_B,$$

so that $u \cdot 1_B = \underbrace{-a_0}_{\in I_1} \cdot 1_B \in I_1 \cdot 1_B$. This proves Lemma \mathcal{G} .]

[*Proof of Lemma \mathcal{H}* : Assume that $u \cdot 1_B \in I_1 \cdot 1_B$. Thus, $u \cdot 1_B = w \cdot 1_B$ for some $w \in I_1$. Consider this w . Then, $w \in I_1$, so that $-w \in -I_1 \subseteq I_1$ (since I_1 is an ideal of A). Define a 2-tuple $(a_0, a_1) \in A^2$ by setting $a_0 = -w$ and $a_1 = 1$. Then,

$$\begin{aligned} \sum_{k=0}^1 a_k (u \cdot 1_B)^k &= \underbrace{a_0}_{=-w} \underbrace{(u \cdot 1_B)^0}_{=1_B} + \underbrace{a_1}_{=1} \underbrace{(u \cdot 1_B)^1}_{=u \cdot 1_B} = -w \cdot 1_B + \underbrace{u \cdot 1_B}_{=w \cdot 1_B} \\ &= -w \cdot 1_B + w \cdot 1_B = 0. \end{aligned}$$

Also, $a_i \in I_{1-i}$ for every $i \in \{0, 1\}$ (since $a_0 = -w \in I_1 = I_{1-0}$ and $a_1 = 1 \in A = I_0 = I_{1-1}$). Altogether, we now know that $(a_0, a_1) \in A^2$ and

$$\sum_{k=0}^1 a_k (u \cdot 1_B)^k = 0, \quad a_1 = 1, \quad \text{and} \quad a_i \in I_{1-i} \text{ for every } i \in \{0, 1\}.$$

Thus, by Definition 2.3 (applied to $u \cdot 1_B$ and 1 instead of u and n), the element $u \cdot 1_B$ is 1-integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. This proves Lemma \mathcal{H} .]

Combining Lemma \mathcal{G} and Lemma \mathcal{H} , we obtain that $u \cdot 1_B$ is 1-integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$ if and only if $u \cdot 1_B \in I_1 \cdot 1_B$. This proves Theorem 2.12. \square

The next theorem is an analogue of Theorem 1.7 (a) for integrality over ideal semifiltrations:

Theorem 2.13. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $x \in B$ and $y \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Assume that x is m -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$, and that y is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$.
Then, $x + y$ is nm -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$.

Proof of Theorem 2.13. Consider the polynomial ring $A[Y]$ and its A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. The polynomial ring $B[Y]$ is an $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ -algebra (as explained in Remark 2.9).

Theorem 2.11 (applied to x and m instead of u and n) yields that xY is m -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ (since x is m -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$). Also, Theorem 2.11 (applied to y instead of u) yields that yY is n -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ (since y is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$).

Hence, Theorem 1.7 (a) (applied to $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$, $B[Y]$, xY and yY instead of A , B , x and y , respectively) yields that $xY + yY$ is nm -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. Since $xY + yY = (x + y)Y$, this means that $(x + y)Y$ is nm -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. Hence, Theorem 2.11 (applied to $x + y$ and nm

instead of u and n) yields that $x + y$ is nm -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. This proves Theorem 2.13. \square

Our next theorem is a somewhat asymmetric analogue of Theorem 1.7 (b) for integrality over ideal semifiltrations:

Theorem 2.14. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $x \in B$ and $y \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Assume that x is m -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$, and that y is n -integral over A .

Then, xy is nm -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$.

Before we prove this theorem, we require a trivial observation:

Lemma 2.15. Let A be a ring. Let A' be an A -algebra. Let B' be an A' -algebra. Let $v \in B'$. Let $n \in \mathbb{N}$. Assume that v is n -integral over A . (Here, of course, we are using the fact that B' is an A -algebra, since B' is an A' -algebra while A' is an A -algebra.)

Then, v is n -integral over A' .

Proof of Lemma 2.15. We know that v is n -integral over A . In other words, there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(v) = 0$. Consider this P , and denote it by Q . Thus, Q is a monic polynomial in $A[X]$ with $\deg Q = n$ and $Q(v) = 0$.

Consider the canonical ring homomorphism $A \rightarrow A'$ sending each $a \in A$ to $a \cdot 1_{A'} \in A'$. This homomorphism is defined because A' is an A -algebra, and it in turn induces a canonical ring homomorphism $A[X] \rightarrow A'[X]$. Let $\tilde{Q} \in A'[X]$ be the image of the polynomial $Q \in A[X]$ under this latter homomorphism. Then, \tilde{Q} is a monic polynomial with $\deg \tilde{Q} = n$ (since Q is a monic polynomial with $\deg Q = n$). Furthermore, the definition of \tilde{Q} yields $\tilde{Q}(v) = Q(v) = 0$. Thus, there exists a monic polynomial $P \in A'[X]$ with $\deg P = n$ and $P(v) = 0$ (namely, $P = \tilde{Q}$). In other words, v is n -integral over A' . This proves Lemma 2.15. \square

Proof of Theorem 2.14. Consider the polynomial ring $A[Y]$ and its A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. The polynomial ring $B[Y]$ is an $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ -algebra (as explained in Remark 2.9).

Theorem 2.11 (applied to x and m instead of u and n) yields that xY is m -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ (since x is m -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$). Also, we know that y is n -integral over A . Thus, Lemma 2.15 (applied to $A' = A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$, $B' = B[Y]$ and $v = y$) yields that y is n -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ (since $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ is an A -algebra, and $B[Y]$ is an $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ -algebra).

On the other hand, we know that xY is m -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Hence, Theorem 1.7 (b) (applied to $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$, $B[Y]$ and xY instead of A , B and x , respectively) yields that $xY \cdot y$ is nm -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Since $xY \cdot y = xyY$, this means that xyY is nm -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Hence, Theorem 2.11 (applied to xy and nm instead of u and n) yields that xy is nm -integral over $\left(A, (I_\rho)_{\rho \in \mathbb{N}} \right)$. This proves Theorem 2.14. \square

It is easy to state analogues of Corollary 1.9 and Corollary 1.10 for ideal semifiltrations. These analogues can be derived from Corollary 1.9 and Corollary 1.10 in the same way as how we derived Theorem 2.13 from Theorem 1.7 (a).

2.5. Transitivity again

The next theorem imitates Theorem 1.5 for integrality over ideal semifiltrations:

Theorem 2.16. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A .

Let $v \in B$ and $u \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$.

(a) Then, $(I_\rho A[v])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[v]$. (See Convention 2.17 below for the meaning of “ $I_\rho A[v]$ ”.)

(b) Assume that v is m -integral over A , and that u is n -integral over $\left(A[v], (I_\rho A[v])_{\rho \in \mathbb{N}} \right)$. Then, u is nm -integral over $\left(A, (I_\rho)_{\rho \in \mathbb{N}} \right)$.

Here and in the following, we are using the following convention:

Convention 2.17. Let A be a ring. Let B be an A -algebra. Let $v \in B$, and let I be an ideal of A . Then, you should read the expression “ $IA[v]$ ” as $I \cdot (A[v])$, not as $(IA)[v]$. For instance, you should read the term “ $I_\rho A[v]$ ” (in Theorem 2.16 (a)) as $I_\rho \cdot (A[v])$, not as $(I_\rho A)[v]$.

Before we prove Theorem 2.16, let us state two lemmas. The first is a more general (but still obvious) version of Theorem 2.16 (a):

Lemma 2.18. Let A be a ring. Let A' be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Then, $(I_\rho A')_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A' .

Proof of Lemma 2.18. We know that $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A . In other words, $(I_\rho)_{\rho \in \mathbb{N}}$ is a sequence of ideals of A and satisfies

$$\begin{aligned} I_0 &= A; \\ I_a I_b &\subseteq I_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N} \end{aligned}$$

(by Definition 2.1). The set I_ρ is an ideal of A for every $\rho \in \mathbb{N}$ (since $(I_\rho)_{\rho \in \mathbb{N}}$ is a sequence of ideals of A).

Now, the set $I_\rho A'$ is an ideal of A' for every $\rho \in \mathbb{N}$ (since I_ρ is an ideal of A). Hence, $(I_\rho A')_{\rho \in \mathbb{N}}$ is a sequence of ideals of A' . It satisfies

$$\begin{aligned} \underbrace{I_0}_{=A} A' &= AA' = A' && \text{(since } A' \text{ is an } A\text{-algebra)}; \\ I_a A' \cdot I_b A' &= \underbrace{I_a I_b}_{\subseteq I_{a+b}} A' \subseteq I_{a+b} A' && \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

Thus, by Definition 2.1 (applied to A' and $(I_\rho A')_{\rho \in \mathbb{N}}$ instead of A and $(I_\rho)_{\rho \in \mathbb{N}}$), it follows that $(I_\rho A')_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A' . This proves Lemma 2.18. \square

Lemma 2.19. Let A be a ring. Let A' be an A -algebra. Let B' be an A' -algebra. Let $v \in B'$. Then, $A' \cdot A[v] = A'[v]$ (an equality between A -submodules of B'). (Here, we are using the fact that B' is an A -algebra, because B' is an A' -algebra while A' is an A -algebra.)

Here, of course, the expression “ $A' \cdot A[v]$ ” means “ $A' \cdot (A[v])$ ”, not “ $(A' \cdot A)[v]$ ”.

Proof of Lemma 2.19. We have $A[v] \subseteq A'[v]$ (since the ring A acts on B' through the canonical ring homomorphism $A \rightarrow A'$). Hence, $A' \cdot A[v] \subseteq A' \cdot \underbrace{A[v]}_{\subseteq A'[v]} \subseteq$

$A'[v]$ (since $A'[v]$ is an A' -algebra). On the other hand, let x be an element of $A'[v]$. Then, there exist some $n \in \mathbb{N}$ and some $(a_0, a_1, \dots, a_n) \in (A')^{n+1}$ such that $x = \sum_{k=0}^n a_k v^k$. Consider this n and this (a_0, a_1, \dots, a_n) . Thus,

$$x = \sum_{k=0}^n \underbrace{a_k}_{\in A'} \underbrace{v^k}_{\in A[v]} \in \sum_{k=0}^n A' \cdot A[v] \subseteq A' \cdot A[v] \quad \text{(since } A' \cdot A[v] \text{ is an additive group)}.$$

Now, forget that we fixed x . Thus, we have proved that $x \in A' \cdot A[v]$ for every $x \in A'[v]$. Therefore, $A'[v] \subseteq A' \cdot A[v]$. Combined with $A' \cdot A[v] \subseteq A'[v]$, this yields $A' \cdot A[v] = A'[v]$. Hence, we have established Lemma 2.19. \square

We are now ready to prove Theorem 2.16:

Proof of Theorem 2.16. **(a)** Lemma 2.18 (applied to $A' = A[v]$) yields that $(I_\rho A[v])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[v]$. This proves Theorem 2.16 **(a)**.

(b) Consider the polynomial ring $A[Y]$ and its A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. Then, $(A[v])[Y]$ is an $A[Y]$ -algebra (since $A[v]$ is an A -algebra) and therefore also an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra (since $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is a subring of $A[Y]$).

Hence, $\left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]\right) [v]$ is an A -subalgebra of $(A[v])[Y]$ (since $v \in A[v] \subseteq (A[v])[Y]$). On the other hand, $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right]$ is an A -subalgebra of $(A[v])[Y]$ (by its definition).

Note that B is an $A[v]$ -algebra (since $A[v]$ is a subring of B). Hence, (as explained in Definition 2.6) the polynomial ring $B[Y]$ is an $(A[v])[Y]$ -algebra. Moreover, $B[Y]$ is an $A[Y]$ -algebra (as explained in Definition 2.6) and also an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra (as explained in Remark 2.9).

Now, we will show that $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right] = \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]\right) [v]$. (This is an equality between two subrings of $(A[v])[Y]$.)

In fact, Definition 2.7 yields $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] = \sum_{i \in \mathbb{N}} I_i Y^i$. The same definition (but applied to $A[v]$ and $(I_\rho A[v])_{\rho \in \mathbb{N}}$ instead of A and $(I_\rho)_{\rho \in \mathbb{N}}$) yields

$$\begin{aligned} (A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right] &= \sum_{i \in \mathbb{N}} I_i \underbrace{A[v] \cdot Y^i}_{=Y^i \cdot A[v]} = \sum_{i \in \mathbb{N}} I_i Y^i \cdot A[v] \\ &= \underbrace{\left(\sum_{i \in \mathbb{N}} I_i Y^i \right)}_{=A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]} \cdot A[v] = A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \cdot A[v] \\ &= \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]\right) [v] \end{aligned} \quad (14)$$

(by Lemma 2.19, applied to $A' = A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ and $B' = (A[v])[Y]$).

Recall that $B[Y]$ is an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra. Hence, Lemma 2.15 (applied to $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$, $B[Y]$ and m instead of A' , B' and n) yields that v is m -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ (since v is m -integral over A).

Now, Theorem 2.11 (applied to $A[v]$ and $(I_\rho A[v])_{\rho \in \mathbb{N}}$ instead of A and $(I_\rho)_{\rho \in \mathbb{N}}$) yields that the element uY is n -integral over $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right]$ (since u is n -integral over $(A[v], (I_\rho A[v])_{\rho \in \mathbb{N}})$). In view of (14), this rewrites as follows: The element uY is n -integral over $\left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]\right) [v]$. Hence, Theorem 1.5 (applied to $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$, $B[Y]$ and uY instead of A , B and u) yields that uY is nm -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ (since v is m -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$). Thus, Theorem 2.11 (applied to nm instead of n) yields that u is nm -integral over $\left(A, (I_\rho)_{\rho \in \mathbb{N}}\right)$. This proves Theorem 2.16 (b). \square

3. Generalizing to two ideal semifiltrations

Theorem 2.14 can be generalized: Instead of requiring y to be integral over the ring A , we can require y to be integral over a further ideal semifiltration $(J_\rho)_{\rho \in \mathbb{N}}$ of A . In that case, xy will be integral over the ideal semifiltration $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ (see Theorem 3.4 for the precise statement). To get a grip on this, let us study two ideal semifiltrations.

3.1. The product of two ideal semifiltrations

Theorem 3.1. Let A be a ring.

(a) Then, $(A)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A .

(b) Let $(I_\rho)_{\rho \in \mathbb{N}}$ and $(J_\rho)_{\rho \in \mathbb{N}}$ be two ideal semifiltrations of A . Then, $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A .

Proof of Theorem 3.1. (a) Clearly, $(A)_{\rho \in \mathbb{N}}$ is a sequence of ideals of A . Hence, in order to prove that $(A)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , it is enough to verify that it satisfies the two conditions

$$\begin{aligned} A &= A; \\ AA &\subseteq A \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

But these two conditions are obviously satisfied. Hence, $(A)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (by Definition 2.1, applied to $(A)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$). This proves Theorem 3.1 (a).

(b) Since $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , it is a sequence of ideals of A , and it satisfies the two conditions

$$\begin{aligned} I_0 &= A; \\ I_a I_b &\subseteq I_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N} \end{aligned} \tag{15}$$

(by Definition 2.1). Since $(J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , it is a sequence of ideals of A , and it satisfies the two conditions

$$\begin{aligned} J_0 &= A; \\ J_a J_b &\subseteq J_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N} \end{aligned} \tag{16}$$

(by Definition 2.1, applied to $(J_\rho)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$).

Now, we know that both $(I_\rho)_{\rho \in \mathbb{N}}$ and $(J_\rho)_{\rho \in \mathbb{N}}$ are sequences of ideals of A . Hence, if $\rho \in \mathbb{N}$, then both I_ρ and J_ρ are ideals of A , and therefore $I_\rho J_\rho$ is an ideal of A as well (since the product of any two ideals of A is an ideal of A). Thus, $I_\rho J_\rho$ is an ideal of A for each $\rho \in \mathbb{N}$. In other words, $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ is a sequence of

ideals of A . Thus, in order to prove that $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , it is enough to verify that it satisfies the two conditions

$$\begin{aligned} I_0 J_0 &= A; \\ I_a J_a \cdot I_b J_b &\subseteq I_{a+b} J_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

But these two conditions are satisfied, since

$$\begin{aligned} \underbrace{I_0}_{=A} \underbrace{J_0}_{=A} &= AA = A; \\ I_a J_a \cdot I_b J_b &= \underbrace{I_a I_b}_{\subseteq I_{a+b} \text{ (by (15))}} \underbrace{J_a J_b}_{\subseteq J_{a+b} \text{ (by (16))}} \subseteq I_{a+b} J_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

Hence, $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (by Definition 2.1, applied to $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$). This proves Theorem 3.1 (b). \square

3.2. Half-reduction

Now let us generalize Theorem 2.11:

Theorem 3.2. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ and $(J_\rho)_{\rho \in \mathbb{N}}$ be two ideal semifiltrations of A . Let $n \in \mathbb{N}$. Let $u \in B$.

We know that $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (according to Theorem 3.1 (b)).

Consider the polynomial ring $A[Y]$ and its A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$.

We will abbreviate this A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ by $A_{[I]}$.

(a) The sequence $\left(J_\tau A_{[I]} \right)_{\tau \in \mathbb{N}}$ is an ideal semifiltration of $A_{[I]}$.

(b) The element u of B is n -integral over $\left(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}} \right)$ if and only if the element uY of the polynomial ring $B[Y]$ is n -integral over $\left(A_{[I]}, \left(J_\tau A_{[I]} \right)_{\tau \in \mathbb{N}} \right)$. (Here, we are using the fact that $B[Y]$ is an $A_{[I]}$ -algebra, because $A_{[I]} = A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is a subring of $A[Y]$ and because $B[Y]$ is an $A[Y]$ -algebra as explained in Definition 2.6.)

Proof of Theorem 3.2. The definition of $A_{[I]}$ yields

$$\begin{aligned} A_{[I]} &= A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] = \sum_{i \in \mathbb{N}} I_i Y^i \quad \text{(by Definition 2.7)} \\ &= \sum_{\ell \in \mathbb{N}} I_\ell Y^\ell \quad \text{(here we renamed } i \text{ as } \ell \text{ in the sum)}. \end{aligned}$$

As a consequence of this chain of equalities, we have $\sum_{i \in \mathbb{N}} I_i Y^i = A_{[I]}$ and $\sum_{\ell \in \mathbb{N}} I_\ell Y^\ell = A_{[I]}$.

(a) We know that $(J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A . In other words, $(J_\tau)_{\tau \in \mathbb{N}}$ is an ideal semifiltration of A (since $(J_\tau)_{\tau \in \mathbb{N}} = (J_\rho)_{\rho \in \mathbb{N}}$). Thus, by Lemma 2.18 (applied to $A_{[I]}$ and $(J_\tau)_{\tau \in \mathbb{N}}$ instead of A' and $(I_\rho)_{\rho \in \mathbb{N}}$), the sequence $(J_\tau A_{[I]})_{\tau \in \mathbb{N}}$ is an ideal semifiltration of $A_{[I]}$. This proves Theorem 3.2 (a).

(b) In order to verify Theorem 3.2 (b), we have to prove the following two lemmata:

Lemma \mathcal{E}' : If u is n -integral over $(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}})$, then uY is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$.

Lemma \mathcal{F}' : If uY is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$, then u is n -integral over $(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}})$.

[Proof of Lemma \mathcal{E}' : Assume that u is n -integral over $(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$), there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} J_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (a_0, a_1, \dots, a_n) .

For every $k \in \{0, 1, \dots, n\}$, we have

$$\begin{aligned} a_k &\in I_{n-k} \underbrace{J_{n-k}}_{\subseteq A} && (\text{since } a_i \in I_{n-i} J_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}) \\ &\subseteq I_{n-k} A \subseteq I_{n-k} && (\text{since } I_{n-k} \text{ is an ideal of } A) \end{aligned}$$

and thus

$$\underbrace{a_k}_{\in I_{n-k}} Y^{n-k} \in I_{n-k} Y^{n-k} \subseteq \sum_{i \in \mathbb{N}} I_i Y^i = A_{[I]}.$$

Thus, we can define an $(n+1)$ -tuple $(b_0, b_1, \dots, b_n) \in (A_{[I]})^{n+1}$ by setting

$$(b_k = a_k Y^{n-k} \text{ for every } k \in \{0, 1, \dots, n\}).$$

Consider this $(n + 1)$ -tuple. The definition of this $(n + 1)$ -tuple yields

$$\sum_{k=0}^n \underbrace{b_k}_{=a_k Y^{n-k}} \cdot \underbrace{(uY)^k}_{=u^k Y^k} = \sum_{k=0}^n a_k Y^{n-k} u^k Y^k = \sum_{k=0}^n a_k u^k \underbrace{Y^{n-k} Y^k}_{=Y^n} = Y^n \cdot \underbrace{\sum_{k=0}^n a_k u^k}_{=0} = 0;$$

$$b_n = \underbrace{a_n}_{=1} \underbrace{Y^{n-n}}_{=Y^0=1} = 1,$$

and

$$b_i = \underbrace{a_i}_{\substack{\in I_{n-i} J_{n-i} \\ = J_{n-i} I_{n-i}}} Y^{n-i} \in J_{n-i} \underbrace{I_{n-i} Y^{n-i}}_{\substack{\subseteq \sum_{\ell \in \mathbb{N}} I_\ell Y^\ell \\ = A_{[I]}}} \subseteq J_{n-i} A_{[I]} \quad \text{for every } i \in \{0, 1, \dots, n\}.$$

Altogether, we now know that $(b_0, b_1, \dots, b_n) \in (A_{[I]})^{n+1}$ and

$$\sum_{k=0}^n b_k \cdot (uY)^k = 0, \quad b_n = 1, \quad \text{and} \quad b_i \in J_{n-i} A_{[I]} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Hence, by Definition 2.3 (applied to $A_{[I]}$, $B[Y]$, $(J_\tau A_{[I]})_{\tau \in \mathbb{N}}$, uY and (b_0, b_1, \dots, b_n) instead of A , B , $(I_\rho)_{\rho \in \mathbb{N}}$, u and (a_0, a_1, \dots, a_n)), the element uY is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. This proves Lemma \mathcal{E}' .]

[Proof of Lemma \mathcal{F}' : Assume that uY is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $A_{[I]}$, $B[Y]$, $(J_\tau A_{[I]})_{\tau \in \mathbb{N}}$, uY and (p_0, p_1, \dots, p_n) instead of A , B , $(I_\rho)_{\rho \in \mathbb{N}}$, u and (a_0, a_1, \dots, a_n)), there exists some $(p_0, p_1, \dots, p_n) \in (A_{[I]})^{n+1}$ such that

$$\sum_{k=0}^n p_k \cdot (uY)^k = 0, \quad p_n = 1, \quad \text{and} \quad p_i \in J_{n-i} A_{[I]} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (p_0, p_1, \dots, p_n) . For every $k \in \{0, 1, \dots, n\}$, we have

$$\begin{aligned} p_k &\in J_{n-k} A_{[I]} && \left(\text{since } p_i \in J_{n-i} A_{[I]} \text{ for every } i \in \{0, 1, \dots, n\} \right) \\ &= J_{n-k} \sum_{i \in \mathbb{N}} I_i Y^i && \left(\text{since } A_{[I]} = \sum_{i \in \mathbb{N}} I_i Y^i \right) \\ &= \sum_{i \in \mathbb{N}} \underbrace{J_{n-k} I_i}_{=I_i J_{n-k}} Y^i = \sum_{i \in \mathbb{N}} I_i J_{n-k} Y^i, \end{aligned}$$

and thus there exists a sequence $(p_{k,i})_{i \in \mathbb{N}} \in A^{\mathbb{N}}$ such that $p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i$, such that $(p_{k,i} \in I_i J_{n-k}$ for every $i \in \mathbb{N}$), and such that only finitely many $i \in \mathbb{N}$ satisfy $p_{k,i} \neq 0$. Consider this sequence. Thus,

$$\begin{aligned}
 & \sum_{k=0}^n \underbrace{p_k}_{= \sum_{i \in \mathbb{N}} p_{k,i} Y^i} \cdot \underbrace{(uY)^k}_{= u^k Y^k = Y^k u^k} \\
 &= \sum_{k=0}^n \left(\sum_{i \in \mathbb{N}} p_{k,i} Y^i \right) \cdot Y^k u^k = \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} \underbrace{Y^i \cdot Y^k}_{= Y^{i+k}} u^k = \sum_{k \in \{0,1,\dots,n\}} \sum_{i \in \mathbb{N}} p_{k,i} Y^{i+k} u^k \\
 &= \sum_{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}} p_{k,i} Y^{i+k} u^k = \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} \underbrace{Y^{i+k}}_{= Y^\ell} u^k \\
 &= \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} Y^\ell u^k = \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} u^k Y^\ell.
 \end{aligned}$$

Comparing this with $\sum_{k=0}^n p_k \cdot (uY)^k = 0$, we obtain $\sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} u^k Y^\ell = 0$.

In other words, the polynomial $\underbrace{\sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=\ell}} p_{k,i} u^k Y^\ell}_{\in B} \in B[Y]$ equals 0.

Hence, its coefficient before Y^n equals 0 as well. But its coefficient before Y^n is $\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=n}} p_{k,i} u^k$. Comparing the preceding two sentences, we see that

$\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=n}} p_{k,i} u^k$ equals 0. Thus,

$$0 = \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+k=n}} p_{k,i} u^k = \sum_{k \in \{0,1,\dots,n\}} \sum_{\substack{i \in \mathbb{N}; \\ i+k=n}} p_{k,i} u^k. \quad (17)$$

But for any given $k \in \{0, 1, \dots, n\}$, we have

$$\left\{ i \in \mathbb{N} \mid \underbrace{i+k=n}_{\Leftrightarrow (i=n-k)} \right\} = \{i \in \mathbb{N} \mid i = n - k\} = \{n - k\}$$

(since $n - k \in \mathbb{N}$ (because $k \in \{0, 1, \dots, n\}$)) and therefore

$$\sum_{\substack{i \in \mathbb{N}; \\ i+k=n}} p_{k,i} u^k = \sum_{i \in \{n-k\}} p_{k,i} u^k = p_{k,n-k} u^k.$$

Hence, (17) becomes

$$0 = \sum_{k \in \{0,1,\dots,n\}} \underbrace{\sum_{\substack{i \in \mathbb{N}; \\ i+k=n}} p_{k,i} u^k}_{=p_{k,n-k}u^k} = \sum_{k \in \{0,1,\dots,n\}} p_{k,n-k} u^k. \quad (18)$$

Recall that $p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i$ for every $k \in \{0,1,\dots,n\}$. Applying this to $k = n$, we find $p_n = \sum_{i \in \mathbb{N}} p_{n,i} Y^i$. Comparing this with $p_n = 1 = 1 \cdot Y^0$, we find

$$\sum_{i \in \mathbb{N}} p_{n,i} Y^i = 1 \cdot Y^0 \quad \text{in } A[Y].$$

Hence, the coefficient of the polynomial $\sum_{i \in \mathbb{N}} p_{n,i} Y^i \in A[Y]$ before Y^0 is 1. But the coefficient of the polynomial $\sum_{i \in \mathbb{N}} p_{n,i} Y^i \in A[Y]$ before Y^0 is $p_{n,0}$ (since $p_{n,i} \in A$ for all $i \in \mathbb{N}$). Comparing the preceding two sentences, we see that $p_{n,0} = 1$.

Define an $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ by setting

$$(a_k = p_{k,n-k} \text{ for every } k \in \{0,1,\dots,n\}).$$

Then, $a_n = p_{n,n-n} = p_{n,0} = 1$. Besides,

$$\sum_{k=0}^n \underbrace{a_k}_{\substack{=p_{k,n-k} \\ \text{(by the definition} \\ \text{of } a_k)}} u^k = \sum_{k=0}^n p_{k,n-k} u^k = \sum_{k \in \{0,1,\dots,n\}} p_{k,n-k} u^k = 0 \quad (\text{by (18)}).$$

Finally, for every $k \in \{0,1,\dots,n\}$, we have $n-k \in \mathbb{N}$ and thus $a_k = p_{k,n-k} \in I_{n-k} J_{n-k}$ (since $p_{k,i} \in I_i J_{n-k}$ for every $i \in \mathbb{N}$). Renaming the variable k as i in this statement, we obtain the following: For every $i \in \{0,1,\dots,n\}$, we have $a_i \in I_{n-i} J_{n-i}$.

Altogether, we now know that the $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ satisfies

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} J_{n-i} \text{ for every } i \in \{0,1,\dots,n\}.$$

Thus, by Definition 2.3 (applied to $(I_\rho J_\rho)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$), the element u is n -integral over $(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}})$. This proves Lemma \mathcal{F}' .]

Combining Lemma \mathcal{E}' and Lemma \mathcal{F}' , we obtain that u is n -integral over $(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}})$ if and only if uY is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. This proves Theorem 3.2 (b). \square

The reason why Theorem 3.2 **(b)** generalizes Theorem 2.11 (more precisely, Theorem 2.11 is the particular case of Theorem 3.2 **(b)** for $J_\rho = A$) is the following fact, which we mention here for the pure sake of completeness:

Theorem 3.3. Let A be a ring. Let B be an A -algebra. Let $n \in \mathbb{N}$. Let $u \in B$.

We know that $(A)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (according to Theorem 3.1 **(a)**).

Then, the element u of B is n -integral over $(A, (A)_{\rho \in \mathbb{N}})$ if and only if u is n -integral over A .

Proof of Theorem 3.3. In order to verify Theorem 3.3, we have to prove the following two lemmata:

Lemma \mathcal{L} : If u is n -integral over $(A, (A)_{\rho \in \mathbb{N}})$, then u is n -integral over A .

Lemma \mathcal{M} : If u is n -integral over A , then u is n -integral over $(A, (A)_{\rho \in \mathbb{N}})$.

[*Proof of Lemma \mathcal{L} :* Assume that u is n -integral over $(A, (A)_{\rho \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $(A)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$), there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in A \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (a_0, a_1, \dots, a_n) .

Define a polynomial $P \in A[X]$ by $P(X) = \sum_{k=0}^n a_k X^k$. Then, $P(X) = \sum_{k=0}^n a_k X^k = \underbrace{a_n}_{=1} X^n + \sum_{k=0}^{n-1} a_k X^k = X^n + \sum_{k=0}^{n-1} a_k X^k$. Hence, the polynomial P is monic, and $\deg P = n$. Besides, $P(u) = 0$ (since $P(X) = \sum_{k=0}^n a_k X^k$ yields $P(u) = \sum_{k=0}^n a_k u^k = 0$). Thus, there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$. Hence, u is n -integral over A . This proves Lemma \mathcal{L} .]

[*Proof of Lemma \mathcal{M} :* Assume that u is n -integral over A . Thus, there exists a monic polynomial $P \in A[X]$ with $\deg P = n$ and $P(u) = 0$. Consider this P . Since $\deg P = n$, there exists some $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that $P(X) = \sum_{k=0}^n a_k X^k$. Consider this (a_0, a_1, \dots, a_n) . Thus, $a_n = 1$ (since P is monic, and $\deg P = n$). Also, substituting u for X in the equality $\sum_{k=0}^n a_k X^k = P(X)$ yields

$\sum_{k=0}^n a_k u^k = P(u) = 0$. Altogether, we now know that $(a_0, a_1, \dots, a_n) \in A^{n+1}$ and

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in A \text{ for every } i \in \{0, 1, \dots, n\}.$$

Hence, by Definition 2.3 (applied to $(A)_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$), the element u is n -integral over $(A, (A)_{\rho \in \mathbb{N}})$. This proves Lemma \mathcal{M} .]

Combining Lemma \mathcal{L} and Lemma \mathcal{M} , we obtain that u is n -integral over $(A, (A)_{\rho \in \mathbb{N}})$ if and only if u is n -integral over A . This proves Theorem 3.3. \square

3.3. Integrality of products over the product semifiltration

Finally, let us generalize Theorem 2.14:

Theorem 3.4. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ and $(J_\rho)_{\rho \in \mathbb{N}}$ be two ideal semifiltrations of A .

Let $x \in B$ and $y \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Assume that x is m -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$, and that y is n -integral over $(A, (J_\rho)_{\rho \in \mathbb{N}})$. Then, xy is nm -integral over $(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}})$.

The proof of this theorem will require a generalization of Lemma 2.15:

Lemma 3.5. Let A be a ring. Let A' be an A -algebra. Let B' be an A' -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $v \in B'$. Let $n \in \mathbb{N}$. Assume that v is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. (Here, of course, we are using the fact that B' is an A -algebra, since B' is an A' -algebra while A' is an A -algebra.)

Then, v is n -integral over $(A', (I_\rho A')_{\rho \in \mathbb{N}})$. (Note that $(I_\rho A')_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A' , according to Lemma 2.18.)

Proof of Lemma 3.5. We know that v is n -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $B = B'$ and $u = v$), there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k v^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (a_0, a_1, \dots, a_n) .

Now, recall that A' is an A -algebra. Define an $(n+1)$ -tuple $(b_0, b_1, \dots, b_n) \in (A')^{n+1}$ by setting

$$(b_i = a_i \cdot 1_{A'} \quad \text{for each } i \in \{0, 1, \dots, n\}).$$

Then, we have $b_i = \underbrace{a_i}_{\in I_{n-i}} \cdot \underbrace{1_{A'}}_{\in A'} \in I_{n-i}A'$ for every $i \in \{0, 1, \dots, n\}$. Also,

$$\sum_{k=0}^n \underbrace{b_k}_{=a_k \cdot 1_{A'}} \quad v^k = \sum_{k=0}^n \underbrace{(a_k \cdot 1_{A'})}_{=a_k v^k} v^k = \sum_{k=0}^n a_k v^k = 0.$$

(by the definition of b_k)

Furthermore, the definition of b_n yields $b_n = \underbrace{a_n}_{=1} \cdot 1_{A'} = 1_{A'} = 1$.

Thus, $(b_0, b_1, \dots, b_n) \in (A')^{n+1}$ and

$$\sum_{k=0}^n b_k v^k = 0, \quad b_n = 1, \quad \text{and} \quad b_i \in I_{n-i}A' \text{ for every } i \in \{0, 1, \dots, n\}.$$

Hence, by Definition 2.3 (applied to B' , A' , $(I_\rho A')_{\rho \in \mathbb{N}}$, v and (b_0, b_1, \dots, b_n) instead of B , A , $(I_\rho)_{\rho \in \mathbb{N}}$, u and (a_0, a_1, \dots, a_n)), the element v is n -integral over $(A', (I_\rho A')_{\rho \in \mathbb{N}})$. This proves Lemma 3.5. \square

Proof of Theorem 3.4. We have $(J_\rho)_{\rho \in \mathbb{N}} = (J_\tau)_{\tau \in \mathbb{N}}$. Hence, y is n -integral over $(A, (J_\tau)_{\tau \in \mathbb{N}})$ (since y is n -integral over $(A, (J_\rho)_{\rho \in \mathbb{N}})$). Also, $(J_\tau)_{\tau \in \mathbb{N}}$ is an ideal semifiltration of A (since $(J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , but we have $(J_\rho)_{\rho \in \mathbb{N}} = (J_\tau)_{\tau \in \mathbb{N}}$). Thus, $(J_\tau A_{[I]})_{\tau \in \mathbb{N}}$ is an ideal semifiltration of $A_{[I]}$ (by Lemma 2.18, applied to $A_{[I]}$ and $(J_\tau)_{\tau \in \mathbb{N}}$ instead of A' and $(I_\rho)_{\rho \in \mathbb{N}}$).

Consider the polynomial ring $A[Y]$ and its A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. We will abbreviate this A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ by $A_{[I]}$. Thus, $A_{[I]} = A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ is a subring of $A[Y]$. Hence, $B[Y]$ is an $A_{[I]}$ -algebra (since $B[Y]$ is an $A[Y]$ -algebra as explained in Definition 2.6).

Theorem 2.11 (applied to x and m instead of u and n) yields that xY is m -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ (since x is m -integral over $(A, (I_\rho)_{\rho \in \mathbb{N}})$). In other words, xY is m -integral over $A_{[I]}$ (since $A[(I_\rho)_{\rho \in \mathbb{N}} * Y] = A_{[I]}$).

On the other hand, $A_{[I]}$ is an A -algebra, and $B[Y]$ is an $A_{[I]}$ -algebra. Hence, Lemma 3.5 (applied to $A_{[I]}$, $B[Y]$, $(J_\tau)_{\tau \in \mathbb{N}}$ and y instead of A' , B' , $(I_\rho)_{\rho \in \mathbb{N}}$ and v) yields that y is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$ (since y is n -integral over $(A, (J_\tau)_{\tau \in \mathbb{N}})$).

Hence, Theorem 2.14 (applied to $A_{[I]}$, $B[Y]$, $(J_\tau A_{[I]})_{\tau \in \mathbb{N}}$, y , xY , n and m instead of A , B , $(I_\rho)_{\rho \in \mathbb{N}}$, x , y , m and n , respectively) yields that $y \cdot xY$ is mn -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$ (since xY is m -integral over $A_{[I]}$).

Since $y \cdot xY = xyY$ and $mn = nm$, this means that xyY is nm -integral over $\left(A_{[I]}, \left(J_\tau A_{[I]}\right)_{\tau \in \mathbb{N}}\right)$. Hence, Theorem 3.2 (b) (applied to xy and nm instead of u and n) yields that xy is nm -integral over $\left(A, (I_\rho J_\rho)_{\rho \in \mathbb{N}}\right)$. This proves Theorem 3.4. \square

4. Accelerating ideal semifiltrations

4.1. Definition of λ -acceleration

We start this section with an obvious observation:

Theorem 4.1. Let A be a ring. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $\lambda \in \mathbb{N}$. Then, $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A .

Proof of Theorem 4.1. Since $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , it is a sequence of ideals of A , and it satisfies the two conditions

$$\begin{aligned} I_0 &= A; \\ I_a I_b &\subseteq I_{a+b} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N} \end{aligned} \tag{19}$$

(by Definition 2.1).

Now, $I_{\lambda\rho}$ is an ideal of A for every $\rho \in \mathbb{N}$ (since $(I_\rho)_{\rho \in \mathbb{N}}$ is a sequence of ideals of A). Hence, $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is a sequence of ideals of A . Thus, in order to prove that $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , it is enough to verify that it satisfies the two conditions

$$\begin{aligned} I_{\lambda \cdot 0} &= A; \\ I_{\lambda a} I_{\lambda b} &\subseteq I_{\lambda(a+b)} \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

But these two conditions are satisfied, since

$$\begin{aligned} I_{\lambda \cdot 0} &= I_0 = A; \\ I_{\lambda a} I_{\lambda b} &\subseteq I_{\lambda a + \lambda b} \quad (\text{by (19), applied to } \lambda a \text{ and } \lambda b \text{ instead of } a \text{ and } b) \\ &= I_{\lambda(a+b)} \quad (\text{since } \lambda a + \lambda b = \lambda(a+b)) \quad \text{for every } a \in \mathbb{N} \text{ and } b \in \mathbb{N}. \end{aligned}$$

Hence, $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (by Definition 2.1, applied to $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$). This proves Theorem 4.1. \square

I refer to the ideal semifiltration $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ in Theorem 4.1 as the λ -acceleration of the ideal semifiltration $(I_\rho)_{\rho \in \mathbb{N}}$.

4.2. Half-reduction and reduction

Now, Theorem 3.2, itself a generalization of Theorem 2.11, can be generalized once more:

Theorem 4.2. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ and $(J_\rho)_{\rho \in \mathbb{N}}$ be two ideal semifiltrations of A . Let $n \in \mathbb{N}$. Let $u \in B$. Let $\lambda \in \mathbb{N}$.

We know that $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (according to Theorem 4.1).

Hence, $(I_{\lambda\rho}J_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (according to Theorem 3.1 (b), applied to $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ instead of $(I_\rho)_{\rho \in \mathbb{N}}$).

Consider the polynomial ring $A[Y]$ and its A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$.

We will abbreviate this A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ by $A_{[I]}$.

(a) The sequence $(J_\tau A_{[I]})_{\tau \in \mathbb{N}}$ is an ideal semifiltration of $A_{[I]}$.

(b) The element u of B is n -integral over $(A, (I_{\lambda\rho}J_\rho)_{\rho \in \mathbb{N}})$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. (Here, we are using the fact that $B[Y]$ is an $A_{[I]}$ -algebra, because $A_{[I]} = A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ is a subring of $A[Y]$ and because $B[Y]$ is an $A[Y]$ -algebra as explained in Definition 2.6.)

Proof of Theorem 4.2. (a) This is precisely the claim of Theorem 3.2 (a); thus, we don't need to prove it again.

(b) The definition of $A_{[I]}$ yields

$$\begin{aligned} A_{[I]} &= A[(I_\rho)_{\rho \in \mathbb{N}} * Y] = \sum_{i \in \mathbb{N}} I_i Y^i && \text{(by Definition 2.7)} \\ &= \sum_{\ell \in \mathbb{N}} I_\ell Y^\ell && \text{(here we renamed } i \text{ as } \ell \text{ in the sum).} \end{aligned}$$

As a consequence of this chain of equalities, we have $\sum_{i \in \mathbb{N}} I_i Y^i = A_{[I]}$ and $\sum_{\ell \in \mathbb{N}} I_\ell Y^\ell = A_{[I]}$.

In order to verify Theorem 4.2 (b), we have to prove the following two lemmata:

Lemma \mathcal{E}'' : If u is n -integral over $(A, (I_{\lambda\rho}J_\rho)_{\rho \in \mathbb{N}})$, then uY^λ is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$.

Lemma \mathcal{F}'' : If uY^λ is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$, then u is n -integral over $(A, (I_{\lambda\rho}J_\rho)_{\rho \in \mathbb{N}})$.

[Proof of Lemma \mathcal{E}'' : Assume that u is n -integral over $(A, (I_{\lambda\rho}J_{\rho})_{\rho \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $(I_{\lambda\rho}J_{\rho})_{\rho \in \mathbb{N}}$ instead of $(I_{\rho})_{\rho \in \mathbb{N}}$), there exists some $(a_0, a_1, \dots, a_n) \in A^{n+1}$ such that

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{\lambda(n-i)}J_{n-i} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (a_0, a_1, \dots, a_n) .

For each $k \in \{0, 1, \dots, n\}$, we have

$$\begin{aligned} a_k &\in I_{\lambda(n-k)} \underbrace{J_{n-k}}_{\subseteq A} && \left(\text{since } a_i \in I_{\lambda(n-i)}J_{n-i} \text{ for every } i \in \{0, 1, \dots, n\} \right) \\ &\subseteq I_{\lambda(n-k)}A \subseteq I_{\lambda(n-k)} && \left(\text{since } I_{\lambda(n-k)} \text{ is an ideal of } A \right) \end{aligned}$$

and thus

$$\underbrace{a_k}_{\in I_{\lambda(n-k)}} Y^{\lambda(n-k)} \in I_{\lambda(n-k)} Y^{\lambda(n-k)} \subseteq \sum_{i \in \mathbb{N}} I_i Y^i = A_{[I]}.$$

Thus, we can define an $(n+1)$ -tuple $(b_0, b_1, \dots, b_n) \in (A_{[I]})^{n+1}$ by

$$(b_k = a_k Y^{\lambda(n-k)} \text{ for every } k \in \{0, 1, \dots, n\}).$$

Consider this $(n+1)$ -tuple. Then,

$$\begin{aligned} \sum_{k=0}^n \underbrace{b_k}_{=a_k Y^{\lambda(n-k)} \text{ (by the definition of } b_k)} \cdot \underbrace{(uY^{\lambda})^k}_{=u^k(Y^{\lambda})^k = u^k Y^{\lambda k}} &= \sum_{k=0}^n a_k \underbrace{Y^{\lambda(n-k)} u^k}_{=u^k Y^{\lambda(n-k)}} Y^{\lambda k} = \sum_{k=0}^n a_k u^k \underbrace{Y^{\lambda(n-k)} Y^{\lambda k}}_{=Y^{\lambda(n-k)+\lambda k} = Y^{\lambda n}} \\ &= \sum_{k=0}^n a_k u^k Y^{\lambda n} = Y^{\lambda n} \cdot \underbrace{\sum_{k=0}^n a_k u^k}_{=0} = 0, \end{aligned}$$

Furthermore, the definition of b_n yields

$$b_n = \underbrace{a_n}_{=1} \underbrace{Y^{\lambda(n-n)}}_{=Y^{\lambda \cdot 0} = Y^0 = 1} = 1.$$

Finally, the definition of b_i yields

$$b_i = \underbrace{a_i}_{\substack{\in I_{\lambda(n-i)}J_{n-i} \\ =J_{n-i}I_{\lambda(n-i)}}} Y^{\lambda(n-i)} \in J_{n-i} \underbrace{I_{\lambda(n-i)} Y^{\lambda(n-i)}}_{\substack{\subseteq \sum_{\ell \in \mathbb{N}} I_{\ell} Y^{\ell} \\ =A_{[I]}}} \subseteq J_{n-i} A_{[I]} \quad \text{for every } i \in \{0, 1, \dots, n\}.$$

Altogether, we now know that $(b_0, b_1, \dots, b_n) \in (A_{[I]})^{n+1}$ and

$$\sum_{k=0}^n b_k \cdot (uY^\lambda)^k = 0, \quad b_n = 1, \quad \text{and} \quad b_i \in J_{n-i}A_{[I]} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Hence, by Definition 2.3 (applied to $A_{[I]}, B[Y], (J_\tau A_{[I]})_{\tau \in \mathbb{N}}, uY^\lambda$ and (b_0, b_1, \dots, b_n) instead of $A, B, (I_\rho)_{\rho \in \mathbb{N}}, u$ and (a_0, a_1, \dots, a_n)), the element uY^λ is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. This proves Lemma \mathcal{E}'' .]

[*Proof of Lemma \mathcal{F}''* : Assume that uY^λ is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. Thus, by Definition 2.3 (applied to $A_{[I]}, B[Y], (J_\tau A_{[I]})_{\tau \in \mathbb{N}}, uY^\lambda$ and (p_0, p_1, \dots, p_n) instead of $A, B, (I_\rho)_{\rho \in \mathbb{N}}, u$ and (a_0, a_1, \dots, a_n)), there exists some $(p_0, p_1, \dots, p_n) \in (A_{[I]})^{n+1}$ such that

$$\sum_{k=0}^n p_k \cdot (uY^\lambda)^k = 0, \quad p_n = 1, \quad \text{and} \quad p_i \in J_{n-i}A_{[I]} \text{ for every } i \in \{0, 1, \dots, n\}.$$

Consider this (p_0, p_1, \dots, p_n) . For every $k \in \{0, 1, \dots, n\}$, we have

$$\begin{aligned} p_k &\in J_{n-k}A_{[I]} && \left(\text{since } p_i \in J_{n-i}A_{[I]} \text{ for every } i \in \{0, 1, \dots, n\} \right) \\ &= J_{n-k} \sum_{i \in \mathbb{N}} I_i Y^i && \left(\text{since } A_{[I]} = \sum_{i \in \mathbb{N}} I_i Y^i \right) \\ &= \sum_{i \in \mathbb{N}} \underbrace{J_{n-k} I_i}_{= I_i J_{n-k}} Y^i = \sum_{i \in \mathbb{N}} I_i J_{n-k} Y^i, \end{aligned}$$

and thus there exists a sequence $(p_{k,i})_{i \in \mathbb{N}} \in A^{\mathbb{N}}$ such that $p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i$, such that $(p_{k,i} \in I_i J_{n-k} \text{ for every } i \in \mathbb{N})$, and such that only finitely many $i \in \mathbb{N}$ sat-

isfy $p_{k,i} \neq 0$. Consider this sequence. Thus,

$$\begin{aligned}
 \sum_{k=0}^n \underbrace{p_k}_{=\sum_{i \in \mathbb{N}} p_{k,i} Y^i} \cdot \underbrace{(uY^\lambda)^k}_{=\underbrace{u^k (\gamma^\lambda)^k}_{=u^k \gamma^{\lambda k}}}_{=Y^{\lambda k} u^k} &= \sum_{k=0}^n \left(\sum_{i \in \mathbb{N}} p_{k,i} Y^i \right) \cdot Y^{\lambda k} u^k \quad \left(\text{since } p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i \right) \\
 &= \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} \underbrace{Y^i \cdot Y^{\lambda k}}_{=Y^{i+\lambda k}} u^k = \sum_{k=0}^n \sum_{i \in \mathbb{N}} p_{k,i} Y^{i+\lambda k} u^k \\
 &= \sum_{k \in \{0,1,\dots,n\}} \sum_{i \in \mathbb{N}} p_{k,i} Y^{i+\lambda k} u^k = \sum_{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}} p_{k,i} Y^{i+\lambda k} u^k \\
 &= \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \ell}} p_{k,i} \underbrace{Y^{i+\lambda k}}_{=Y^\ell} u^k \quad (\text{since } i+\lambda k = \ell) \\
 &= \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \ell}} p_{k,i} \underbrace{Y^\ell u^k}_{=u^k Y^\ell} = \sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \ell}} p_{k,i} u^k Y^\ell.
 \end{aligned}$$

Comparing this with $\sum_{k=0}^n p_k \cdot (uY^\lambda)^k = 0$, we obtain $\sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \ell}} p_{k,i} u^k Y^\ell =$

0. In other words, the polynomial $\underbrace{\sum_{\ell \in \mathbb{N}} \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \ell}} p_{k,i} u^k Y^\ell}_{\in B} \in B[Y]$ equals 0.

Hence, its coefficient before $Y^{\lambda n}$ equals 0 as well. But its coefficient before $Y^{\lambda n}$ is $\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \lambda n}} p_{k,i} u^k$. Comparing the preceding two sentences, we see that

$\sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \lambda n}} p_{k,i} u^k$ equals 0. Thus,

$$0 = \sum_{\substack{(k,i) \in \{0,1,\dots,n\} \times \mathbb{N}; \\ i+\lambda k = \lambda n}} p_{k,i} u^k = \sum_{k \in \{0,1,\dots,n\}} \sum_{\substack{i \in \mathbb{N}; \\ i+\lambda k = \lambda n}} p_{k,i} u^k. \quad (20)$$

But for each given $k \in \{0,1,\dots,n\}$, we have $n-k \in \mathbb{N}$ and thus $\lambda(n-k) \in \mathbb{N}$ (since $\lambda \in \mathbb{N}$) and thus

$$\begin{aligned}
 \left\{ i \in \mathbb{N} \mid \underbrace{i + \lambda k = \lambda n}_{\iff (i = \lambda n - \lambda k)} \right\} &= \left\{ i \in \mathbb{N} \mid i = \underbrace{\lambda n - \lambda k}_{= \lambda(n-k)} \right\} \\
 &= \{ i \in \mathbb{N} \mid i = \lambda(n-k) \} = \{ \lambda(n-k) \}
 \end{aligned}$$

(since $\lambda(n-k) \in \mathbb{N}$) and therefore

$$\sum_{\substack{i \in \mathbb{N}; \\ i + \lambda k = \lambda n}} p_{k,i} u^k = \sum_{i \in \{\lambda(n-k)\}} p_{k,i} u^k = p_{k,\lambda(n-k)} u^k.$$

Hence, (20) becomes

$$0 = \sum_{k \in \{0,1,\dots,n\}} \underbrace{\sum_{\substack{i \in \mathbb{N}; \\ i + \lambda k = \lambda n}} p_{k,i} u^k}_{= p_{k,\lambda(n-k)} u^k} = \sum_{k \in \{0,1,\dots,n\}} p_{k,\lambda(n-k)} u^k. \quad (21)$$

Recall that $p_k = \sum_{i \in \mathbb{N}} p_{k,i} Y^i$ for every $k \in \{0,1,\dots,n\}$. Applying this to $k = n$, we find $p_n = \sum_{i \in \mathbb{N}} p_{n,i} Y^i$. Comparing this with $p_n = 1 = 1 \cdot Y^0$, we find

$$\sum_{i \in \mathbb{N}} p_{n,i} Y^i = 1 \cdot Y^0 \quad \text{in } A[Y].$$

Hence, the coefficient of the polynomial $\sum_{i \in \mathbb{N}} p_{n,i} Y^i \in A[Y]$ before Y^0 is 1. But the coefficient of the polynomial $\sum_{i \in \mathbb{N}} p_{n,i} Y^i \in A[Y]$ before Y^0 is $p_{n,0}$ (since $p_{n,i} \in A$ for all $i \in \mathbb{N}$). Comparing the preceding two sentences, we see that $p_{n,0} = 1$.

Define an $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ by setting

$$\left(a_k = p_{k,\lambda(n-k)} \text{ for every } k \in \{0,1,\dots,n\} \right).$$

Then, $a_n = p_{n,\lambda(n-n)} = p_{n,\lambda \cdot 0} = p_{n,0} = 1$. Besides,

$$\sum_{k=0}^n \underbrace{a_k}_{\substack{= p_{k,\lambda(n-k)} \\ \text{(by the definition} \\ \text{of } a_k)}} u^k = \sum_{k=0}^n p_{k,\lambda(n-k)} u^k = \sum_{k \in \{0,1,\dots,n\}} p_{k,\lambda(n-k)} u^k = 0 \quad (\text{by (21)}).$$

Finally, for every $k \in \{0,1,\dots,n\}$, we have $n-k \in \mathbb{N}$ and therefore $\lambda(n-k) \in \mathbb{N}$ (since $\lambda \in \mathbb{N}$) and thus $a_k = p_{k,\lambda(n-k)} \in I_{\lambda(n-k)} J_{n-k}$ (since $p_{k,i} \in I_i J_{n-k}$ for every $i \in \mathbb{N}$). Renaming the variable k as i in this statement, we obtain the following: For every $i \in \{0,1,\dots,n\}$, we have $a_i \in I_{\lambda(n-i)} J_{n-i}$.

Altogether, we now know that the $(n+1)$ -tuple $(a_0, a_1, \dots, a_n) \in A^{n+1}$ satisfies

$$\sum_{k=0}^n a_k u^k = 0, \quad a_n = 1, \quad \text{and} \quad a_i \in I_{\lambda(n-i)} J_{n-i} \text{ for every } i \in \{0,1,\dots,n\}.$$

Thus, by Definition 2.3 (applied to $(I_{\lambda\rho} J_{\rho})_{\rho \in \mathbb{N}}$ instead of $(I_{\rho})_{\rho \in \mathbb{N}}$), the element u is n -integral over $\left(A, (I_{\lambda\rho} J_{\rho})_{\rho \in \mathbb{N}} \right)$. This proves Lemma \mathcal{F}'' .]

Combining Lemma \mathcal{E}'' and Lemma \mathcal{F}'' , we obtain that u is n -integral over $(A, (I_{\lambda\rho}J\rho)_{\rho \in \mathbb{N}})$ if and only if uY^λ is n -integral over $(A_{[I]}, (J_\tau A_{[I]})_{\tau \in \mathbb{N}})$. This proves Theorem 4.2 (b). \square

A particular case of Theorem 4.2 (b) is the following fact:

Theorem 4.3. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $n \in \mathbb{N}$. Let $u \in B$. Let $\lambda \in \mathbb{N}$.

We know that $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (according to Theorem 4.1).

Consider the polynomial ring $A[Y]$ and its A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ defined in Definition 2.7.

Then, the element u of B is n -integral over $(A, (I_{\lambda\rho})_{\rho \in \mathbb{N}})$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over the ring $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. (Here, we are using the fact that $B[Y]$ is an $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ -algebra, because $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ is a subring of $A[Y]$ and because $B[Y]$ is an $A[Y]$ -algebra as explained in Definition 2.6.)

Proof of Theorem 4.3. Theorem 3.1 (a) states that $(A)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A .

We have $(I_{\lambda\rho})_{\rho \in \mathbb{N}} = (I_{\lambda\rho}A)_{\rho \in \mathbb{N}}$ ⁴.

We will abbreviate the A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ of $A[Y]$ by $A_{[I]}$. Thus, $B[Y]$ is an $A_{[I]}$ -algebra (since $B[Y]$ is an $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ -algebra).

It is easy to see that $AA_{[I]} = A_{[I]}$ ⁵. Hence, $\left(\underbrace{AA_{[I]}}_{=A_{[I]}} \right)_{\tau \in \mathbb{N}} = (A_{[I]})_{\tau \in \mathbb{N}} = (A_{[I]})_{\rho \in \mathbb{N}}$.

We have the following five equivalences:

⁴*Proof.* We know that $(I_{\lambda\rho})_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , thus a sequence of ideals of A . In other words, for each $\rho \in \mathbb{N}$, the set $I_{\lambda\rho}$ is an ideal of A .

Now, let $\rho \in \mathbb{N}$. Then, the set $I_{\lambda\rho}$ is an ideal of A (as we have just seen). Hence, $I_{\lambda\rho}A \subseteq I_{\lambda\rho}$. Combining this with $I_{\lambda\rho} = I_{\lambda\rho} \underbrace{1_A}_{\in A} \subseteq I_{\lambda\rho}A$, we obtain $I_{\lambda\rho} = I_{\lambda\rho}A$.

Forget that we fixed ρ . We thus have shown that $I_{\lambda\rho} = I_{\lambda\rho}A$ for each $\rho \in \mathbb{N}$. In other words, $(I_{\lambda\rho})_{\rho \in \mathbb{N}} = (I_{\lambda\rho}A)_{\rho \in \mathbb{N}}$.

⁵*Proof.* We have $AA_{[I]} \subseteq A_{[I]}$ (since $A_{[I]}$ is an A -algebra). Combining this with $A_{[I]} = \underbrace{1_A}_{\in A} \cdot A_{[I]} \subseteq AA_{[I]}$, we obtain $AA_{[I]} = A_{[I]}$, qed.

- The element u of B is n -integral over $\left(A, (I_{\lambda\rho})_{\rho \in \mathbb{N}}\right)$ if and only if the element u of B is n -integral over $\left(A, (I_{\lambda\rho}A)_{\rho \in \mathbb{N}}\right)$ (since $(I_{\lambda\rho})_{\rho \in \mathbb{N}} = (I_{\lambda\rho}A)_{\rho \in \mathbb{N}}$).
- The element u of B is n -integral over $\left(A, (I_{\lambda\rho}A)_{\rho \in \mathbb{N}}\right)$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over $\left(A_{[I]}, \left(AA_{[I]}\right)_{\tau \in \mathbb{N}}\right)$ (according to Theorem 4.2 (b), applied to $(A)_{\rho \in \mathbb{N}}$ instead of $(J_\rho)_{\rho \in \mathbb{N}}$).
- The element uY^λ of the polynomial ring $B[Y]$ is n -integral over $\left(A_{[I]}, \left(AA_{[I]}\right)_{\tau \in \mathbb{N}}\right)$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over $\left(A_{[I]}, \left(A_{[I]}\right)_{\rho \in \mathbb{N}}\right)$ (since $\left(AA_{[I]}\right)_{\tau \in \mathbb{N}} = \left(A_{[I]}\right)_{\rho \in \mathbb{N}}$).
- The element uY^λ of the polynomial ring $B[Y]$ is n -integral over $\left(A_{[I]}, \left(A_{[I]}\right)_{\rho \in \mathbb{N}}\right)$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over $A_{[I]}$ (by Theorem 3.3, applied to $A_{[I]}, B[Y]$ and uY^λ instead of A, B and u).
- The element uY^λ of the polynomial ring $B[Y]$ is n -integral over $A_{[I]}$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ (since $A_{[I]} = A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$).

Combining these five equivalences, we obtain that the element u of B is n -integral over $\left(A, (I_{\lambda\rho})_{\rho \in \mathbb{N}}\right)$ if and only if the element uY^λ of the polynomial ring $B[Y]$ is n -integral over $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$. This proves Theorem 4.3. \square

Note that Theorem 2.11 is the particular case of Theorem 4.3 for $\lambda = 1$. Finally we can generalize even Theorem 1.11:

Theorem 4.4. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $n \in \mathbb{N}^+$. Let $v \in B$. Let a_0, a_1, \dots, a_n be $n+1$ elements of A such that $\sum_{i=0}^n a_i v^i = 0$. Assume further that $a_i \in I_{n-i}$ for every $i \in \{0, 1, \dots, n\}$.

Let $k \in \{0, 1, \dots, n\}$. We know that $\left(I_{(n-k)\rho}\right)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A (according to Theorem 4.1, applied to $\lambda = n - k$).

Then, $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -integral over $\left(A, \left(I_{(n-k)\rho}\right)_{\rho \in \mathbb{N}}\right)$.

Proof of Theorem 4.4. Consider the polynomial ring $A[Y]$ and its A -subalgebra $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ defined in Definition 2.7. Note that $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is a subring

of $A[Y]$; hence, $B[Y]$ is an $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ -algebra (because $B[Y]$ is an $A[Y]$ -algebra as explained in Definition 2.6).

Definition 2.7 yields

$$A[(I_\rho)_{\rho \in \mathbb{N}} * Y] = \sum_{i \in \mathbb{N}} I_i Y^i = \sum_{\ell \in \mathbb{N}} I_\ell Y^\ell \quad (\text{here we renamed } i \text{ as } \ell \text{ in the sum}).$$

$$\text{Hence, } \sum_{\ell \in \mathbb{N}} I_\ell Y^\ell = A[(I_\rho)_{\rho \in \mathbb{N}} * Y].$$

Define $u \in B$ by

$$u = \sum_{i=0}^{n-k} a_{i+k} v^i. \quad (22)$$

In the ring $B[Y]$, we have

$$\sum_{i=0}^n a_i Y^{n-i} \underbrace{(vY)^i}_{=v^i Y^i = Y^i v^i} = \sum_{i=0}^n a_i \underbrace{Y^{n-i} Y^i}_{=Y^n} v^i = Y^n \sum_{i=0}^n a_i v^i = 0.$$

Besides, every $i \in \{0, 1, \dots, n\}$ satisfies

$$\underbrace{a_i}_{\substack{\in I_{n-i} \\ \text{(by assumption)}}} Y^{n-i} \in I_{n-i} Y^{n-i} \subseteq \sum_{\ell \in \mathbb{N}} I_\ell Y^\ell = A[(I_\rho)_{\rho \in \mathbb{N}} * Y].$$

In other words, $a_0 Y^{n-0}, a_1 Y^{n-1}, \dots, a_n Y^{n-n}$ are $n+1$ elements of $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$.

Hence, Theorem 1.11 (applied to $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$, $B[Y]$, vY and $a_i Y^{n-i}$ instead of A , B , v and a_i) yields that $\sum_{i=0}^{n-k} a_{i+k} Y^{n-(i+k)} (vY)^i$ is n -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$.

Since

$$\sum_{i=0}^{n-k} a_{i+k} Y^{n-(i+k)} \underbrace{(vY)^i}_{=v^i Y^i = Y^i v^i} = \sum_{i=0}^{n-k} a_{i+k} \underbrace{Y^{n-(i+k)} Y^i}_{=Y^{n-(i+k)+i} = Y^{n-k}} v^i = \underbrace{\sum_{i=0}^{n-k} a_{i+k} v^i}_{=u \text{ (by (22))}} \cdot Y^{n-k} = u Y^{n-k},$$

this means that $u Y^{n-k}$ is n -integral over $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$.

But Theorem 4.3 (applied to $\lambda = n - k$) yields that u is n -integral over

$$\left(A, \left(I_{(n-k)\rho} \right)_{\rho \in \mathbb{N}} \right) \text{ if and only if } u Y^{n-k} \text{ is } n\text{-integral over the ring } A[(I_\rho)_{\rho \in \mathbb{N}} * Y].$$

Since we know that $u Y^{n-k}$ is n -integral over the ring $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$, this yields

that u is n -integral over $\left(A, \left(I_{(n-k)\rho} \right)_{\rho \in \mathbb{N}} \right)$. In other words, $\sum_{i=0}^{n-k} a_{i+k} v^i$ is n -

integral over $\left(A, \left(I_{(n-k)\rho} \right)_{\rho \in \mathbb{N}} \right)$ (since $u = \sum_{i=0}^{n-k} a_{i+k} v^i$). This proves Theorem 4.4. \square

5. On a lemma by Lombardi

5.1. A lemma on products of powers

Now, we shall show a rather technical lemma:

Lemma 5.1. Let A be a ring. Let B be an A -algebra. Let $x \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Let $u \in B$. Let $\mu \in \mathbb{N}$ and $\nu \in \mathbb{N}$ be such that $\mu + \nu \in \mathbb{N}^+$. Assume that

$$u^n \in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^\nu \rangle_A \quad (23)$$

and that

$$\begin{aligned} u^m x^\mu &\in \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^\mu \rangle_A \\ &\quad + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \langle x^0, x^1, \dots, x^{\mu-1} \rangle_A. \end{aligned} \quad (24)$$

Then, u is $(n\mu + m\nu)$ -integral over A .

This lemma can be seen as a variant of [6, Theorem 2]⁶. Indeed, the particular case of [6, Theorem 2] when $J = 0$ can easily be obtained from Lemma 5.1 (applied to x and α instead of u and x).

Before we prove Lemma 5.1, we recall a basic mathematical principle:

Proposition 5.2. Let $\mathfrak{A}(i)$ be an assertion for every $i \in \mathbb{N}$. If

every $I \in \mathbb{N}$ satisfying $(\mathfrak{A}(i) \text{ for every } i \in \mathbb{N} \text{ such that } i < I)$ satisfies $\mathfrak{A}(I)$,

then

$$\text{every } i \in \mathbb{N} \text{ satisfies } \mathfrak{A}(i).$$

Proposition 5.2 is known as the *principle of strong induction*. By renaming i , I and \mathfrak{A} as j , J and \mathfrak{B} , respectively, we can rewrite this principle as follows:

Proposition 5.3. Let $\mathfrak{B}(j)$ be an assertion for every $j \in \mathbb{N}$. If

every $J \in \mathbb{N}$ satisfying $(\mathfrak{B}(j) \text{ for every } j \in \mathbb{N} \text{ such that } j < J)$ satisfies $\mathfrak{B}(J)$,

then

$$\text{every } j \in \mathbb{N} \text{ satisfies } \mathfrak{B}(j).$$

Proof of Lemma 5.1. Define the set

$$\begin{aligned} S &= (\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}) \\ &\quad \cup (\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+\nu-1\}). \end{aligned} \quad (25)$$

⁶*Caveat:* The notion “integral over (A, J) ” defined in [6] has nothing to do with our notion “ n -integral over $(A, (I_n)_{n \in \mathbb{N}}$)”.

Then, $|S| = n\mu + mv$ ⁷. Also,

$$j < \mu + v \text{ for every } (i, j) \in S \quad (26)$$

8.

⁷*Proof.* We have $(U \times V) \cap (X \times Y) = (U \cap X) \times (V \cap Y)$ for any four sets U, V, X and Y . Applying this to $U = \{0, 1, \dots, n-1\}$, $V = \{0, 1, \dots, \mu-1\}$, $X = \{0, 1, \dots, m-1\}$ and $Y = \{\mu, \mu+1, \dots, \mu+v-1\}$, we find

$$\begin{aligned} & (\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}) \cap (\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+v-1\}) \\ &= (\{0, 1, \dots, n-1\} \cap \{0, 1, \dots, m-1\}) \times \underbrace{(\{0, 1, \dots, \mu-1\} \cap \{\mu, \mu+1, \dots, \mu+v-1\})}_{=\emptyset} \\ &= (\{0, 1, \dots, n-1\} \cap \{0, 1, \dots, m-1\}) \times \emptyset = \emptyset. \end{aligned}$$

Hence,

$$\begin{aligned} & |(\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}) \cup (\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+v-1\})| \\ &= \underbrace{|\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}|}_{=|\{0, 1, \dots, n-1\}| \cdot |\{0, 1, \dots, \mu-1\}|} + \underbrace{|\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+v-1\}|}_{=|\{0, 1, \dots, m-1\}| \cdot |\{\mu, \mu+1, \dots, \mu+v-1\}|} \\ & \quad \left(\begin{array}{c} \text{because any two finite sets } U \text{ and } V \text{ satisfying } U \cap V = \emptyset \\ \text{satisfy } |U \cup V| = |U| + |V| \end{array} \right) \\ &= \underbrace{|\{0, 1, \dots, n-1\}|}_{=n} \cdot \underbrace{|\{0, 1, \dots, \mu-1\}|}_{=\mu} + \underbrace{|\{0, 1, \dots, m-1\}|}_{=m} \cdot \underbrace{|\{\mu, \mu+1, \dots, \mu+v-1\}|}_{=v} = n\mu + mv. \end{aligned}$$

In view of

$$S = (\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}) \cup (\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+v-1\}),$$

this rewrites as $|S| = n\mu + mv$.

⁸In fact, $v \geq 0$ (since $v \in \mathbb{N}$), so that $\underbrace{v}_{\geq 0} - 1 \geq \mu - 1$. Hence, $\mu - 1 \leq \mu + v - 1$, so that

$$\begin{aligned} S &= \left(\begin{array}{c} \{0, 1, \dots, n-1\} \times \underbrace{\{0, 1, \dots, \mu-1\}}_{\substack{\subseteq \{0, 1, \dots, \mu+v-1\} \\ (\text{since } \mu-1 \leq \mu+v-1)}} \end{array} \right) \\ & \quad \cup \left(\begin{array}{c} \{0, 1, \dots, m-1\} \times \underbrace{\{\mu, \mu+1, \dots, \mu+v-1\}}_{\substack{\subseteq \{0, 1, \dots, \mu+v-1\} \\ (\text{since } \mu \geq 0)}} \end{array} \right) \\ &\subseteq (\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu+v-1\}) \\ & \quad \cup (\{0, 1, \dots, m-1\} \times \{0, 1, \dots, \mu+v-1\}) \\ &= (\{0, 1, \dots, n-1\} \cup \{0, 1, \dots, m-1\}) \times \{0, 1, \dots, \mu+v-1\} \end{aligned}$$

(since $(U \times X) \cup (V \times X) = (U \cup V) \times X$ for any three sets U, V and X). Hence, for every $(i, j) \in S$, we have $(i, j) \in S \subseteq (\{0, 1, \dots, n-1\} \cup \{0, 1, \dots, m-1\}) \times \{0, 1, \dots, \mu+v-1\}$ and thus $j \in \{0, 1, \dots, \mu+v-1\}$ and thus $j < \mu + v$. This proves (26).

Let U be the A -submodule $\langle u^i x^j \mid (i, j) \in S \rangle_A$ of B . Then, U is an $(n\mu + mv)$ -generated A -module (since $|S| = n\mu + mv$). Besides, clearly,

$$u^i x^j \in U \text{ for every } (i, j) \in S \quad (27)$$

(since $U = \langle u^i x^j \mid (i, j) \in S \rangle_A$).

Now, we will show that

$$\text{every } i \in \mathbb{N} \text{ and } j \in \mathbb{N} \text{ satisfying } j < \mu + \nu \text{ satisfy } u^i x^j \in U. \quad (28)$$

[*Proof of (28)*]. For every $i \in \mathbb{N}$, define an assertion $\mathfrak{A}(i)$ by

$$\mathfrak{A}(i) = \left(\text{every } j \in \mathbb{N} \text{ satisfies } \left(\text{if } j < \mu + \nu, \text{ then } u^i x^j \in U \right) \right).$$

Let us now show that

$$\text{every } I \in \mathbb{N} \text{ satisfying } (\mathfrak{A}(i) \text{ for every } i \in \mathbb{N} \text{ such that } i < I) \text{ satisfies } \mathfrak{A}(I). \quad (29)$$

[*Proof of (29)*]. Let $I \in \mathbb{N}$ be such that

$$(\mathfrak{A}(i) \text{ for every } i \in \mathbb{N} \text{ such that } i < I). \quad (30)$$

We must prove that $\mathfrak{A}(I)$ holds.

The definition of the assertion $\mathfrak{A}(I)$ yields

$$\mathfrak{A}(I) = \left(\text{every } j \in \mathbb{N} \text{ satisfies } \left(\text{if } j < \mu + \nu, \text{ then } u^I x^j \in U \right) \right).$$

For every $j \in \mathbb{N}$, define an assertion $\mathfrak{B}(j)$ by

$$\mathfrak{B}(j) = \left(\text{if } j < \mu + \nu, \text{ then } u^I x^j \in U \right). \quad (31)$$

Let us now show that

$$\text{every } J \in \mathbb{N} \text{ satisfying } (\mathfrak{B}(j) \text{ for every } j \in \mathbb{N} \text{ such that } j < J) \text{ satisfies } \mathfrak{B}(J). \quad (32)$$

[*Proof of (32)*]. Let $J \in \mathbb{N}$ be such that

$$(\mathfrak{B}(j) \text{ for every } j \in \mathbb{N} \text{ such that } j < J). \quad (33)$$

We must prove that $\mathfrak{B}(J)$ holds.

The definition of the assertion $\mathfrak{B}(J)$ yields

$$\mathfrak{B}(J) = \left(\text{if } J < \mu + \nu, \text{ then } u^I x^J \in U \right).$$

Assume that $J < \mu + \nu$. Then, for every $j \in \mathbb{N}$ such that $j < J$, the assertion $\mathfrak{B}(j)$ holds (due to (33)). In other words, for every $j \in \mathbb{N}$ such that $j < J$, we

have (if $j < \mu + \nu$, then $u^I x^j \in U$) (because this is precisely what the assertion $\mathfrak{B}(j)$ says) and therefore $u^I x^j \in U$ (since $j < \mu + \nu$ automatically holds⁹). Thus we have shown that

$$u^I x^j \in U \text{ for every } j \in \mathbb{N} \text{ such that } j < J. \quad (34)$$

In other words,

$$u^I x^j \in U \text{ for every } j \in \{0, 1, \dots, J-1\} \quad (35)$$

(since the numbers $j \in \{0, 1, \dots, J-1\}$ are precisely the numbers $j \in \mathbb{N}$ such that $j < J$). Hence,

$$\begin{aligned} \sum_{j \in \{0, 1, \dots, J-1\}} a_j u^I x^j &\in U \\ \text{for every } (a_j)_{j \in \{0, 1, \dots, J-1\}} &\in A^{\{0, 1, \dots, J-1\}} \end{aligned} \quad (36)$$

(since U is an A -module, and thus is closed under A -linear combination).

Also, if $i \in \mathbb{N}$ satisfies $i < I$, then the assertion $\mathfrak{A}(i)$ holds (by (30)). In view of the definition of $\mathfrak{A}(i)$, we can restate this as follows: If $i \in \mathbb{N}$ satisfies $i < I$, then every $j \in \mathbb{N}$ satisfies (if $j < \mu + \nu$, then $u^i x^j \in U$). In other words,

$$u^i x^j \in U \text{ for every } i \in \mathbb{N} \text{ and } j \in \mathbb{N} \text{ such that } i < I \text{ and } j < \mu + \nu. \quad (37)$$

Hence,

$$u^i x^j \in U \text{ for every } (i, j) \in \{0, 1, \dots, I-1\} \times \{0, 1, \dots, \mu + \nu - 1\} \quad (38)$$

(because for every $(i, j) \in \{0, 1, \dots, I-1\} \times \{0, 1, \dots, \mu + \nu - 1\}$, we have $i < I$ (since $i \in \{0, 1, \dots, I-1\}$) and $j < \mu + \nu$ (since $j \in \{0, 1, \dots, \mu + \nu - 1\}$) and therefore $u^i x^j \in U$ (by (37))). Hence,

$$\begin{aligned} \sum_{(i, j) \in \{0, 1, \dots, I-1\} \times \{0, 1, \dots, \mu + \nu - 1\}} a_{i, j} u^i x^j &\in U \\ \text{for every } (a_{i, j})_{(i, j) \in \{0, 1, \dots, I-1\} \times \{0, 1, \dots, \mu + \nu - 1\}} &\in A^{\{0, 1, \dots, I-1\} \times \{0, 1, \dots, \mu + \nu - 1\}} \end{aligned} \quad (39)$$

(since U is an A -module, and thus is closed under A -linear combination).

Now,

$$\begin{aligned} &\left\langle u^I \right\rangle_A \cdot \underbrace{\left\langle x^0, x^1, \dots, x^{J-1} \right\rangle_A}_{=\langle x^j \mid j \in \{0, 1, \dots, J-1\} \rangle_A} \\ &= \left\langle u^I \right\rangle_A \cdot \left\langle x^j \mid j \in \{0, 1, \dots, J-1\} \right\rangle_A = \left\langle u^I x^j \mid j \in \{0, 1, \dots, J-1\} \right\rangle_A \\ &= \left\{ \sum_{j \in \{0, 1, \dots, J-1\}} a_j u^I x^j \mid (a_j)_{j \in \{0, 1, \dots, J-1\}} \in A^{\{0, 1, \dots, J-1\}} \right\} \subseteq U \end{aligned} \quad (40)$$

⁹because $j < J < \mu + \nu$

(by (36)).

Furthermore,

$$\begin{aligned}
 & \underbrace{\langle u^0, u^1, \dots, u^{I-1} \rangle_A}_{= \langle u^i \mid i \in \{0, 1, \dots, I-1\} \rangle_A} \cdot \underbrace{\langle x^0, x^1, \dots, x^{\mu+\nu-1} \rangle_A}_{= \langle x^j \mid j \in \{0, 1, \dots, \mu+\nu-1\} \rangle_A} \\
 &= \langle u^i \mid i \in \{0, 1, \dots, I-1\} \rangle_A \cdot \langle x^j \mid j \in \{0, 1, \dots, \mu+\nu-1\} \rangle_A \\
 &= \langle u^i x^j \mid (i, j) \in \{0, 1, \dots, I-1\} \times \{0, 1, \dots, \mu+\nu-1\} \rangle_A \\
 &= \left\{ \begin{array}{l} \sum_{(i,j) \in \{0,1,\dots,I-1\} \times \{0,1,\dots,\mu+\nu-1\}} a_{i,j} u^i x^j \\ \mid (a_{i,j})_{(i,j) \in \{0,1,\dots,I-1\} \times \{0,1,\dots,\mu+\nu-1\}} \in A^{\{0,1,\dots,I-1\} \times \{0,1,\dots,\mu+\nu-1\}} \end{array} \right\} \\
 &\subseteq U
 \end{aligned} \tag{41}$$

(by (39)).

From $J < \mu + \nu$, we obtain $J \leq \mu + \nu - 1$ (since J and $\mu + \nu$ are integers). We are now going to show that $u^I x^J \in U$.

Trivially, we have¹⁰

$$(I \geq m \wedge J \geq \mu) \vee (I < m \wedge J \geq \mu) \vee (I \geq n \wedge J < \mu) \vee (I < n \wedge J < \mu)$$

¹¹. Hence, one of the following four cases must hold:

Case 1: We have $I \geq m \wedge J \geq \mu$.

Case 2: We have $I < m \wedge J \geq \mu$.

Case 3: We have $I \geq n \wedge J < \mu$.

Case 4: We have $I < n \wedge J < \mu$.

Let us first consider Case 1. In this case, we have $I \geq m$ and $J \geq \mu$. Hence,

¹⁰Here, an expression like " $I \geq m \wedge J \geq \mu$ " should be read as " $(I \geq m) \wedge (J \geq \mu)$ ".

¹¹since

$$\begin{aligned}
 & \underbrace{(I \geq m \wedge J \geq \mu) \vee (I < m \wedge J \geq \mu)}_{= (I \geq m \vee I < m) \wedge (J \geq \mu)} \vee \underbrace{(I \geq n \wedge J < \mu) \vee (I < n \wedge J < \mu)}_{= (I \geq n \vee I < n) \wedge (J < \mu)} \\
 & \quad \quad \quad = (J \geq \mu) \quad \quad \quad = (J < \mu) \\
 & \quad \quad \quad \text{(since } (I \geq m \vee I < m) \text{ is true)} \quad \quad \quad \text{(since } (I \geq n \vee I < n) \text{ is true)} \\
 & = (J \geq \mu) \vee (J < \mu) = \text{true}
 \end{aligned}$$

$I - m \geq 0$ (since $I \geq m$) and $J - \mu \geq 0$ (since $J \geq \mu$). Thus,

$$\begin{aligned}
& \underbrace{u^I}_{=u^{I-m}u^m} \underbrace{x^J}_{=x^\mu x^{J-\mu}} \\
& \text{(since } I \geq m \text{) (since } J \geq \mu \text{)} \\
& = u^{I-m} \underbrace{u^m x^\mu}_{\substack{\in \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^\mu \rangle_A + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \langle x^0, x^1, \dots, x^{\mu-1} \rangle_A \\ \text{(by (24))}}} x^{J-\mu} \\
& \in u^{I-m} \left(\left\langle u^0, u^1, \dots, u^{m-1} \right\rangle_A \cdot \left\langle x^0, x^1, \dots, x^\mu \right\rangle_A \right. \\
& \quad \left. + \left\langle u^0, u^1, \dots, u^m \right\rangle_A \cdot \left\langle x^0, x^1, \dots, x^{\mu-1} \right\rangle_A \right) x^{J-\mu} \\
& = \underbrace{u^{I-m} \left\langle u^0, u^1, \dots, u^{m-1} \right\rangle_A}_{= \langle u^{I-m} u^0, u^{I-m} u^1, \dots, u^{I-m} u^{m-1} \rangle_A} \cdot \underbrace{\left\langle x^0, x^1, \dots, x^\mu \right\rangle_A x^{J-\mu}}_{= \langle x^0 x^{J-\mu}, x^1 x^{J-\mu}, \dots, x^\mu x^{J-\mu} \rangle_A} \\
& = \underbrace{\langle u^{(I-m)+0}, u^{(I-m)+1}, \dots, u^{(I-m)+(m-1)} \rangle_A}_{= \langle u^{I-m}, u^{I-m+1}, \dots, u^{I-1} \rangle_A} \cdot \underbrace{\langle x^{0+(J-\mu)}, x^{1+(J-\mu)}, \dots, x^{\mu+(J-\mu)} \rangle_A}_{= \langle x^{J-\mu}, x^{J-\mu+1}, \dots, x^J \rangle_A} \\
& \quad \subseteq \langle u^0, u^1, \dots, u^{I-1} \rangle_A \quad \subseteq \langle x^0, x^1, \dots, x^{\mu+J-1} \rangle_A \\
& \text{(since } \{I-m, I-m+1, \dots, I-1\} \subseteq \{0, 1, \dots, I-1\} \text{ (since } I-m \geq 0 \text{))} \quad \text{(since } \{J-\mu, J-\mu+1, \dots, J\} \subseteq \{0, 1, \dots, \mu+J-1\} \\
& \quad \text{(since } J-\mu \geq 0 \text{ and } J \leq \mu+J-1 \text{))} \\
& + \underbrace{u^{I-m} \left\langle u^0, u^1, \dots, u^m \right\rangle_A}_{= \langle u^{I-m} u^0, u^{I-m} u^1, \dots, u^{I-m} u^m \rangle_A} \cdot \underbrace{\left\langle x^0, x^1, \dots, x^{\mu-1} \right\rangle_A x^{J-\mu}}_{= \langle x^0 x^{J-\mu}, x^1 x^{J-\mu}, \dots, x^{\mu-1} x^{J-\mu} \rangle_A} \\
& = \underbrace{\langle u^{(I-m)+0}, u^{(I-m)+1}, \dots, u^{(I-m)+m} \rangle_A}_{= \langle u^{I-m}, u^{I-m+1}, \dots, u^I \rangle_A} \cdot \underbrace{\langle x^{0+(J-\mu)}, x^{1+(J-\mu)}, \dots, x^{(\mu-1)+(J-\mu)} \rangle_A}_{= \langle x^{J-\mu}, x^{J-\mu+1}, \dots, x^{J-1} \rangle_A} \\
& \quad \subseteq \langle u^0, u^1, \dots, u^I \rangle_A \quad \subseteq \langle x^0, x^1, \dots, x^{J-1} \rangle_A \\
& \text{(since } \{I-m, I-m+1, \dots, I\} \subseteq \{0, 1, \dots, I\} \text{ (since } I-m \geq 0 \text{))} \quad \text{(since } \{J-\mu, J-\mu+1, \dots, J-1\} \subseteq \{0, 1, \dots, J-1\} \\
& \quad \text{(since } J-\mu \geq 0 \text{))}
\end{aligned}$$

$$\begin{aligned}
& \subseteq \underbrace{\langle u^0, u^1, \dots, u^{I-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^{\mu+\nu-1} \rangle_A}_{\substack{\subseteq U \\ \text{(by (41))}}} \\
& \quad + \underbrace{\langle u^0, u^1, \dots, u^I \rangle_A}_{=\langle u^0, u^1, \dots, u^{I-1} \rangle_A + \langle u^I \rangle_A} \cdot \langle x^0, x^1, \dots, x^{J-1} \rangle_A \\
& \subseteq U + \underbrace{\left(\langle u^0, u^1, \dots, u^{I-1} \rangle_A + \langle u^I \rangle_A \right) \cdot \langle x^0, x^1, \dots, x^{J-1} \rangle_A}_{=\langle u^0, u^1, \dots, u^{I-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^{J-1} \rangle_A + \langle u^I \rangle_A \cdot \langle x^0, x^1, \dots, x^{J-1} \rangle_A} \\
& = U + \langle u^0, u^1, \dots, u^{I-1} \rangle_A \cdot \underbrace{\langle x^0, x^1, \dots, x^{J-1} \rangle_A}_{\substack{\subseteq \langle x^0, x^1, \dots, x^{\mu+\nu-1} \rangle_A \\ \text{(since } \{0, 1, \dots, J-1\} \subseteq \{0, 1, \dots, \mu+\nu-1\} \\ \text{(since } J-1 \leq J \leq \mu+\nu-1))}} + \langle u^I \rangle_A \cdot \langle x^0, x^1, \dots, x^{J-1} \rangle_A \\
& \subseteq U + \underbrace{\langle u^0, u^1, \dots, u^{I-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^{\mu+\nu-1} \rangle_A}_{\substack{\subseteq U \\ \text{(by (41))}}} + \underbrace{\langle u^I \rangle_A \cdot \langle x^0, x^1, \dots, x^{J-1} \rangle_A}_{\substack{\subseteq U \\ \text{(by (40))}}} \\
& \subseteq U + U + U \subseteq U \quad (\text{since } U \text{ is an } A\text{-module}).
\end{aligned}$$

Thus, we have proved that $u^I x^J \in U$ holds in Case 1.

Let us next consider Case 2. In this case, we have $I < m$ and $J \geq \mu$. Thus, $I \in \{0, 1, \dots, m-1\}$ (since $I < m$ and $I \in \mathbb{N}$) and $J \in \{\mu, \mu+1, \dots, \mu+\nu-1\}$ (since $J \geq \mu$ and $J < \mu+\nu$). Thus,

$$\begin{aligned}
& (I, J) \in \{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+\nu-1\} \\
& \subseteq (\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}) \\
& \quad \cup (\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+\nu-1\}) \\
& = S \quad (\text{by (25)}).
\end{aligned}$$

Hence, $u^I x^J \in U$ (by (27), applied to I and J instead of i and j). Thus, we have proved that $u^I x^J \in U$ holds in Case 2.

Let us next consider Case 3. In this case, we have $I \geq n$ and $J < \mu$. Hence, $I - n \geq 0$ (since $I \geq n$) and $J + \nu \leq \mu + \nu - 1$ (since $\underbrace{J}_{< \mu} + \nu < \mu + \nu$, and since

$J + \nu$ and $\mu + \nu$ are integers). Thus,

$$\begin{aligned}
 \underbrace{u^I}_{=u^{I-n}u^n \text{ (since } I \geq n)} x^J &= u^{I-n} \underbrace{u^n}_{\substack{\in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \\ \text{(by (23))}}} x^J \\
 &\in \underbrace{u^{I-n} \langle u^0, u^1, \dots, u^{n-1} \rangle_A}_{\substack{= \langle u^{I-n}u^0, u^{I-n}u^1, \dots, u^{I-n}u^{n-1} \rangle_A \\ = \langle u^{(I-n)+0}, u^{(I-n)+1}, \dots, u^{(I-n)+(n-1)} \rangle_A \\ = \langle u^{I-n}, u^{I-n+1}, \dots, u^{I-1} \rangle_A \\ \subseteq \langle u^0, u^1, \dots, u^{I-1} \rangle_A \\ \text{(since } \{I-n, I-n+1, \dots, I-1\} \subseteq \{0, 1, \dots, I-1\} \\ \text{(since } I-n \geq 0))}} \cdot \underbrace{\langle x^0, x^1, \dots, x^\nu \rangle_A x^J}_{\substack{= \langle x^0 x^J, x^1 x^J, \dots, x^\nu x^J \rangle_A \\ = \langle x^{0+J}, x^{1+J}, \dots, x^{\nu+J} \rangle_A \\ = \langle x^J, x^{J+1}, \dots, x^{J+\nu} \rangle_A \\ \subseteq \langle x^0, x^1, \dots, x^{\mu+\nu-1} \rangle_A \\ \text{(since } \{J, J+1, \dots, J+\nu\} \subseteq \{0, 1, \dots, \mu+\nu-1\} \\ \text{(since } J \geq 0 \text{ and } J+\nu \leq \mu+\nu-1))}} \\
 &\subseteq \langle u^0, u^1, \dots, u^{I-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^{\mu+\nu-1} \rangle_A \subseteq U \quad \text{(by (41)).}
 \end{aligned}$$

Thus, we have proved that $u^I x^J \in U$ holds in Case 3.

Finally, let us consider Case 4. In this case, we have $I < n$ and $J < \mu$. Thus, $I \in \{0, 1, \dots, n-1\}$ (since $I < n$ and $I \in \mathbb{N}$) and $J \in \{0, 1, \dots, \mu-1\}$ (since $J < \mu$ and $J \in \mathbb{N}$). Thus,

$$\begin{aligned}
 (I, J) &\in \{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\} \\
 &\subseteq (\{0, 1, \dots, n-1\} \times \{0, 1, \dots, \mu-1\}) \\
 &\quad \cup (\{0, 1, \dots, m-1\} \times \{\mu, \mu+1, \dots, \mu+\nu-1\}) \\
 &= S \quad \text{(by (25)),}
 \end{aligned}$$

so that $u^I x^J \in U$ (by (27), applied to I and J instead of i and j). Thus, we have proved that $u^I x^J \in U$ holds in Case 4.

Therefore, we have proved that $u^I x^J \in U$ holds in each of the four Cases 1, 2, 3 and 4. Hence, $u^I x^J \in U$ always holds.

Now, forget our assumption that $J < \mu + \nu$. Hence, we have proved that if $J < \mu + \nu$, then $u^I x^J \in U$. In other words, we have proved the assertion $\mathfrak{B}(J)$ (because $\mathfrak{B}(J) = (\text{if } J < \mu + \nu, \text{ then } u^I x^J \in U)$).

Thus, we have proved (32).]

Hence, Proposition 5.3 yields that

$$\text{every } j \in \mathbb{N} \text{ satisfies } \mathfrak{B}(j).$$

In other words,

$$\text{every } j \in \mathbb{N} \text{ satisfies } \left(\text{if } j < \mu + \nu, \text{ then } u^I x^j \in U \right)$$

(because of (31)). In other words, the assertion $\mathfrak{A}(I)$ holds (because $\mathfrak{A}(I) = (\text{every } j \in \mathbb{N} \text{ satisfies } (\text{if } j < \mu + \nu, \text{ then } u^I x^j \in U))$).

Thus, we have proved (29).]

Hence, Proposition 5.2 yields that

$$\text{every } i \in \mathbb{N} \text{ satisfies } \mathfrak{A}(i).$$

In other words,

$$\text{every } i \in \mathbb{N} \text{ satisfies } \left(\text{every } j \in \mathbb{N} \text{ satisfies } \left(\text{if } j < \mu + \nu, \text{ then } u^i x^j \in U \right) \right)$$

(since $\mathfrak{A}(i) = (\text{every } j \in \mathbb{N} \text{ satisfies } (\text{if } j < \mu + \nu, \text{ then } u^i x^j \in U))$). This is equivalent to (28). Thus, (28) is proven.]

We have $0 < \mu + \nu$ (since $\mu + \nu \in \mathbb{N}^+$). Thus, we can apply (28) to $i = 0$ and $j = 0$. As a result, we obtain $u^0 x^0 \in U$. In view of $\underbrace{u^0}_{=1} \underbrace{x^0}_{=1} = 1$, this rewrites as

$$1 \in U.$$

Furthermore, if $i \in \mathbb{N}$ and $j \in \mathbb{N}$ satisfy $j < \mu + \nu$, then

$$\underbrace{u \cdot u^i}_{=u^{i+1}} x^j = u^{i+1} x^j \in U$$

(by (28) (applied to $i + 1$ instead of i)). Hence,

$$u \cdot u^i x^j \in U \text{ for every } (i, j) \in S, \tag{42}$$

because every $(i, j) \in S$ satisfies $i \in \mathbb{N}$ and $j \in \mathbb{N}$ and $j < \mu + \nu$ (by (26)). Hence, $\sum_{(i,j) \in S} a_{i,j} \underbrace{u \cdot u^i x^j}_{\substack{\in U \\ \text{(by (42))}}} \in U$ for every $(a_{i,j})_{(i,j) \in S} \in A^S$ (since U is an A -module and thus

is closed under A -linear combination).

Now, from $U = \langle u^i x^j \mid (i, j) \in S \rangle_A$, we obtain

$$\begin{aligned} uU &= u \langle u^i x^j \mid (i, j) \in S \rangle_A = \langle u \cdot u^i x^j \mid (i, j) \in S \rangle_A \\ &= \left\{ \sum_{(i,j) \in S} a_{i,j} u \cdot u^i x^j \mid (a_{i,j})_{(i,j) \in S} \in A^S \right\} \subseteq U \end{aligned}$$

(because $\sum_{(i,j) \in S} a_{i,j} u \cdot u^i x^j \in U$ for every $(a_{i,j})_{(i,j) \in S} \in A^S$).

Altogether, U is an $(n\mu + m\nu)$ -generated A -submodule of B such that $1 \in U$ and $uU \subseteq U$. Thus, $u \in B$ satisfies Assertion \mathcal{C} of Theorem 1.1 with n replaced by $n\mu + m\nu$. Hence, $u \in B$ satisfies the four equivalent assertions \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} of Theorem 1.1 with n replaced by $n\mu + m\nu$. Consequently, u is $(n\mu + m\nu)$ -integral over A . This proves Lemma 5.1. \square

We record a weaker variant of Lemma 5.1:

Lemma 5.4. Let A be a ring. Let B be an A -algebra. Let $x \in B$ and $y \in B$ be such that $xy \in A$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Let $u \in B$. Let $\mu \in \mathbb{N}$ and $\nu \in \mathbb{N}$ be such that $\mu + \nu \in \mathbb{N}^+$. Assume that

$$u^n \in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^\nu \rangle_A \quad (43)$$

and that

$$\begin{aligned} u^m \in & \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle y^0, y^1, \dots, y^\mu \rangle_A \\ & + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \langle y^1, y^2, \dots, y^\mu \rangle_A. \end{aligned} \quad (44)$$

Then, u is $(n\mu + m\nu)$ -integral over A .

Proof of Lemma 5.4. Fix $p \in \mathbb{N}$.

Let $i \in \{p, p+1, \dots, \mu\}$. Thus, $i \geq p$ and $i \leq \mu$. From $i \in \{p, p+1, \dots, \mu\}$, we obtain $\mu - i \in \{0, 1, \dots, \mu - p\}$, so that $\{\mu - i\} \subseteq \{0, 1, \dots, \mu - p\}$. Also, $i \leq \mu$, thus $\mu - i \geq 0$, so that

$$\begin{aligned} y^i \underbrace{x^\mu}_{=x^{\mu-i}x^i} &= y^i x^{\mu-i} x^i = \underbrace{x^i y^i}_{=(xy)^i \in A} x^{\mu-i} \in Ax^{\mu-i} = \langle x^{\mu-i} \rangle_A \end{aligned} \quad (45)$$

(since $xy \in A$)

$$\subseteq \langle x^0, x^1, \dots, x^{\mu-p} \rangle_A \quad (46)$$

(since $\{\mu - i\} \subseteq \{0, 1, \dots, \mu - p\}$).

Now, forget that we fixed i . We thus have proven (46) for each $i \in \{p, p+1, \dots, \mu\}$. Hence, every $(a_i)_{i \in \{p, p+1, \dots, \mu\}} \in A^{\{p, p+1, \dots, \mu\}}$ satisfies

$$\sum_{i \in \{p, p+1, \dots, \mu\}} a_i \underbrace{y^i x^\mu}_{\in \langle x^0, x^1, \dots, x^{\mu-p} \rangle_A} \in \sum_{i \in \{p, p+1, \dots, \mu\}} a_i \langle x^0, x^1, \dots, x^{\mu-p} \rangle_A \subseteq \langle x^0, x^1, \dots, x^{\mu-p} \rangle_A$$

(by (46))

(because $\langle x^0, x^1, \dots, x^{\mu-p} \rangle_A$ is an A -module). In other words,

$$\left\{ \sum_{i \in \{p, p+1, \dots, \mu\}} a_i y^i x^\mu \mid (a_i)_{i \in \{p, p+1, \dots, \mu\}} \in A^{\{p, p+1, \dots, \mu\}} \right\} \subseteq \langle x^0, x^1, \dots, x^{\mu-p} \rangle_A. \quad (47)$$

Now,

$$\begin{aligned}
 \underbrace{\langle y^p, y^{p+1}, \dots, y^\mu \rangle_A}_{= \langle y^i \mid i \in \{p, p+1, \dots, \mu\} \rangle_A} x^\mu &= \langle y^i \mid i \in \{p, p+1, \dots, \mu\} \rangle_A x^\mu \\
 &= \langle y^i x^\mu \mid i \in \{p, p+1, \dots, \mu\} \rangle_A \\
 &= \left\{ \sum_{i \in \{p, p+1, \dots, \mu\}} a_i y^i x^\mu \mid (a_i)_{i \in \{p, p+1, \dots, \mu\}} \in A^{\{p, p+1, \dots, \mu\}} \right\} \\
 &\subseteq \langle x^0, x^1, \dots, x^{\mu-p} \rangle_A \tag{48}
 \end{aligned}$$

(by (47)).

Forget that we fixed p . We thus have proven (48) for each $p \in \mathbb{N}$. Applying (48) to $p = 0$, we find

$$\langle y^0, y^1, \dots, y^\mu \rangle_A x^\mu \subseteq \langle x^0, x^1, \dots, x^{\mu-0} \rangle_A = \langle x^0, x^1, \dots, x^\mu \rangle_A \tag{49}$$

(since $\mu - 0 = \mu$). Applying (48) to $p = 1$, we find

$$\langle y^1, y^2, \dots, y^\mu \rangle_A x^\mu \subseteq \langle x^0, x^1, \dots, x^{\mu-1} \rangle_A. \tag{50}$$

Now, (44) yields

$$\begin{aligned}
 &u^m x^\mu \\
 &\in \left(\langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle y^0, y^1, \dots, y^\mu \rangle_A + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \langle y^1, y^2, \dots, y^\mu \rangle_A \right) x^\mu \\
 &= \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \underbrace{\langle y^0, y^1, \dots, y^\mu \rangle_A x^\mu}_{\substack{\subseteq \langle x^0, x^1, \dots, x^\mu \rangle_A \\ \text{(by (49))}}} + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \underbrace{\langle y^1, y^2, \dots, y^\mu \rangle_A x^\mu}_{\substack{\subseteq \langle x^0, x^1, \dots, x^{\mu-1} \rangle_A \\ \text{(by (50))}}} \\
 &\subseteq \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^\mu \rangle_A + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \langle x^0, x^1, \dots, x^{\mu-1} \rangle_A.
 \end{aligned}$$

In other words, (24) holds. Also, (23) holds (because (43) holds, and because (23) is the same as (43)). Thus, Lemma 5.1 yields that u is $(n\mu + mv)$ -integral over A . This proves Lemma 5.4. \square

We now come to something trivial:

Lemma 5.5. Let A be a ring. Let B be an A -algebra. Let $x \in B$. Let $n \in \mathbb{N}$. Let $u \in B$. Assume that u is n -integral over $A[x]$. Then, there exists some $\nu \in \mathbb{N}^+$ such that

$$u^n \in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^\nu \rangle_A.$$

Proof of Lemma 5.5. There exists a monic polynomial $P \in (A[x])[X]$ with $\deg P = n$ and $P(u) = 0$ (since u is n -integral over $A[x]$). Consider this P . Since $P \in (A[x])[X]$ is a monic polynomial with $\deg P = n$, there exist elements $\alpha_0, \alpha_1, \dots, \alpha_{n-1}$ of $A[x]$ such that $P(X) = X^n + \sum_{i=0}^{n-1} \alpha_i X^i$. Consider these $\alpha_0, \alpha_1, \dots, \alpha_{n-1}$.

Substituting u for X in the equality $P(X) = X^n + \sum_{i=0}^{n-1} \alpha_i X^i$, we find $P(u) = u^n + \sum_{i=0}^{n-1} \alpha_i u^i$. Comparing this with $P(u) = 0$, we obtain $u^n + \sum_{i=0}^{n-1} \alpha_i u^i = 0$.

Hence, $u^n = -\sum_{i=0}^{n-1} \alpha_i u^i$.

For every $i \in \{0, 1, \dots, n-1\}$, we have $\alpha_i \in A[x]$, and thus there exist some $v_i \in \mathbb{N}$ and some $(\beta_{i,0}, \beta_{i,1}, \dots, \beta_{i,v_i}) \in A^{v_i+1}$ such that $\alpha_i = \sum_{k=0}^{v_i} \beta_{i,k} x^k$. Consider these v_i and $(\beta_{i,0}, \beta_{i,1}, \dots, \beta_{i,v_i})$. Hence, for every $i \in \{0, 1, \dots, n-1\}$, we have

$$\alpha_i = \sum_{k=0}^{v_i} \beta_{i,k} x^k \in \langle x^0, x^1, \dots, x^{v_i} \rangle_A. \quad (51)$$

Let $v = \max\{v_0, v_1, \dots, v_{n-1}, 1\}$. Thus, v is an integer satisfying $v \geq 1$ (since $1 \in \{v_0, v_1, \dots, v_{n-1}, 1\}$); hence, $v \in \mathbb{N}^+$. Furthermore, for every $i \in \{0, 1, \dots, n-1\}$, we have $v_i \in \{v_0, v_1, \dots, v_{n-1}\} \subseteq \{v_0, v_1, \dots, v_{n-1}, 1\}$ and thus $v_i \leq \max\{v_0, v_1, \dots, v_{n-1}, 1\} = v$, hence $\{0, 1, \dots, v_i\} \subseteq \{0, 1, \dots, v\}$, and thus

$$\begin{aligned} \alpha_i &\in \langle x^0, x^1, \dots, x^{v_i} \rangle_A && \text{(by (51))} \\ &\subseteq \langle x^0, x^1, \dots, x^v \rangle_A && (52) \end{aligned}$$

(since $\{0, 1, \dots, v_i\} \subseteq \{0, 1, \dots, v\}$). Therefore,

$$\begin{aligned} u^n &= -\sum_{i=0}^{n-1} \alpha_i u^i = -\sum_{i=0}^{n-1} \underbrace{u^i}_{\substack{\in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \\ \text{(since } i \in \{0, 1, \dots, n-1\})}} \cdot \underbrace{\alpha_i}_{\substack{\in \langle x^0, x^1, \dots, x^v \rangle_A \\ \text{(by (52))}}} \\ &\in -\sum_{i=0}^{n-1} \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^v \rangle_A \\ &\subseteq \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^v \rangle_A \end{aligned}$$

(since $\langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^v \rangle_A$ is an A -module). This proves Lemma 5.5. \square

5.2. Integrality over $A[x]$ and over $A[y]$ implies integrality over $A[xy]$

A consequence of Lemma 5.4 and Lemma 5.5 is the following theorem:

Theorem 5.6. Let A be a ring. Let B be an A -algebra. Let $x \in B$ and $y \in B$ be such that $xy \in A$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Let $u \in B$. Assume that u is n -integral over $A[x]$, and that u is m -integral over $A[y]$. Then, there exists some $\lambda \in \mathbb{N}$ such that u is λ -integral over A .

Proof of Theorem 5.6. Since u is n -integral over $A[x]$, Lemma 5.5 yields that there exists some $v \in \mathbb{N}^+$ such that

$$u^n \in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle x^0, x^1, \dots, x^v \rangle_A.$$

In other words, there exists some $v \in \mathbb{N}^+$ such that (43) holds. Consider this v .

Since u is m -integral over $A[y]$, Lemma 5.5 (with x, n and v replaced by y, m and μ) yields that there exists some $\mu \in \mathbb{N}^+$ such that

$$u^m \in \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle y^0, y^1, \dots, y^\mu \rangle_A.$$

Consider this μ . Hence,

$$\begin{aligned} u^m &\in \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle y^0, y^1, \dots, y^\mu \rangle_A \\ &\subseteq \langle u^0, u^1, \dots, u^{m-1} \rangle_A \cdot \langle y^0, y^1, \dots, y^\mu \rangle_A + \langle u^0, u^1, \dots, u^m \rangle_A \cdot \langle y^1, y^2, \dots, y^\mu \rangle_A. \end{aligned}$$

In other words, (44) holds. From $\mu \in \mathbb{N}^+$ and $v \in \mathbb{N}^+$, we obtain $\mu + v \in \mathbb{N}^+$.

Since both (43) and (44) hold, Lemma 5.4 yields that u is $(n\mu + mv)$ -integral over A . Thus, there exists some $\lambda \in \mathbb{N}$ such that u is λ -integral over A (namely, $\lambda = n\mu + mv$). This proves Theorem 5.6. \square

We record a generalization of Theorem 5.6 (which will turn out to be easily seen equivalent to Theorem 5.6):

Theorem 5.7. Let A be a ring. Let B be an A -algebra. Let $x \in B$ and $y \in B$. Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Let $u \in B$. Assume that u is n -integral over $A[x]$, and that u is m -integral over $A[y]$. Then, there exists some $\lambda \in \mathbb{N}$ such that u is λ -integral over $A[xy]$.

Proof of Theorem 5.7. Let C denote the A -subalgebra $A[xy]$ of A . Thus, $C = A[xy]$ is an A -subalgebra of B , hence a subring of B . Thus, $C[x]$ is a C -subalgebra of B , hence a subring of B . Note that $C = A[xy] = A[yx]$ (since $xy = yx$).

Furthermore, $A[x]$ is a subring of $C[x]$ ¹². Thus, $C[x]$ is an $A[x]$ -algebra. Also, B is a $C[x]$ -algebra (since $C[x]$ is a subring of B). Since u is n -integral over $A[x]$, Lemma 2.15 (applied to $B, C[x], A[x]$ and u instead of B', A', A and v)

¹²*Proof.* Both $A[x]$ and $C[x]$ are subrings of B .

Now, let $\gamma \in A[x]$. Thus, there exist some $p \in \mathbb{N}$ and some elements a_0, a_1, \dots, a_p of A such that $\gamma = \sum_{i=0}^p a_i x^i$. Consider this p and these a_0, a_1, \dots, a_p . For each $i \in \{0, 1, \dots, p\}$, we

yields that u is n -integral over $C[x]$. The same argument (but applied to y , x , n and m instead of x , y , m and n) shows that u is m -integral over $C[y]$ (since $C = A[yx]$).

Now, B is a C -algebra (since C is a subring of B) and we have $xy \in A[xy] = C$. Hence, Theorem 5.6 (applied to C instead of A) yields that there exists some $\lambda \in \mathbb{N}$ such that u is λ -integral over C (because u is n -integral over $C[x]$, and because u is m -integral over $C[y]$). In other words, there exists some $\lambda \in \mathbb{N}$ such that u is λ -integral over $A[xy]$ (since $C = A[xy]$). This proves Theorem 5.7. \square

5.3. Generalization to ideal semifiltrations

Theorem 5.7 has a “relative version”:

Theorem 5.8. Let A be a ring. Let B be an A -algebra. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Let $x \in B$ and $y \in B$.

(a) Then, $(I_\rho A[x])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[x]$. Besides, $(I_\rho A[y])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[y]$. Besides, $(I_\rho A[xy])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[xy]$.

(b) Let $m \in \mathbb{N}$ and $n \in \mathbb{N}$. Let $u \in B$. Assume that u is n -integral over $(A[x], (I_\rho A[x])_{\rho \in \mathbb{N}})$, and that u is m -integral over $(A[y], (I_\rho A[y])_{\rho \in \mathbb{N}})$. Then, there exists some $\lambda \in \mathbb{N}$ such that u is λ -integral over $(A[xy], (I_\rho A[xy])_{\rho \in \mathbb{N}})$.

Our proof of this theorem will rely on a lemma:

Lemma 5.9. Let A be a ring. Let B be an A -algebra. Let $v \in B$. Let $(I_\rho)_{\rho \in \mathbb{N}}$ be an ideal semifiltration of A . Lemma 2.18 (applied to $A' = A[v]$) yields that $(I_\rho A[v])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[v]$. Consider the polynomial ring $A[Y]$ and its A -subalgebra $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$. We know that $A[(I_\rho)_{\rho \in \mathbb{N}} * Y]$ is a subring of $A[Y]$, and (as explained in Definition 2.6) the polynomial ring $(A[v])[Y]$ is an $A[Y]$ -algebra (since $A[v]$ is an A -algebra). Hence, $(A[v])[Y]$

have $\underbrace{a_i}_{\in A} \cdot 1_B \in A \cdot 1_B \subseteq A[xy] = C$ (since $C = A[xy]$). Hence, $\sum_{i=0}^p (a_i \cdot 1_B) x^i \in C[x]$. In view of

$$\sum_{i=0}^p (a_i \cdot 1_B) x^i = \sum_{i=0}^p a_i \cdot \underbrace{1_B x^i}_{=x^i} = \sum_{i=0}^p a_i x^i = \gamma \quad \left(\text{since } \gamma = \sum_{i=0}^p a_i x^i \right),$$

this rewrites as $\gamma \in C[x]$.

Forget that we fixed γ . We thus have shown that $\gamma \in C[x]$ for each $\gamma \in A[x]$. In other words, $A[x] \subseteq C[x]$. Hence, $A[x]$ is a subring of $C[x]$ (since both $A[x]$ and $C[x]$ are subrings of B).

is an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra (since $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ is a subring of $A[Y]$). On the other hand, $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right] \subseteq (A[v])[Y]$.

(a) We have

$$(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right] = \left(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right] \right) [v]. \quad (53)$$

(b) Let $u \in B$. Let $n \in \mathbb{N}$. Then, the element u of B is n -integral over $(A[v], (I_\rho A[v])_{\rho \in \mathbb{N}})$ if and only if the element uY of the polynomial ring $B[Y]$ is n -integral over the ring $(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]) [v]$.

Proof of Lemma 5.9. (a) We have proven Lemma 5.9 (a) during the proof of Theorem 2.16 (b).

(b) The ring B is an $A[v]$ -algebra (since $A[v]$ is a subring of B). Hence, Theorem 2.11 (applied to $A[v]$ and $(I_\rho A[v])_{\rho \in \mathbb{N}}$ instead of A and $(I_\rho)_{\rho \in \mathbb{N}}$) yields that the element u of B is n -integral over $(A[v], (I_\rho A[v])_{\rho \in \mathbb{N}})$ if and only if the element uY of the polynomial ring $B[Y]$ is n -integral over the ring $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right]$. In view of (53), this rewrites as follows: The element u of B is n -integral over $(A[v], (I_\rho A[v])_{\rho \in \mathbb{N}})$ if and only if the element uY of the polynomial ring $B[Y]$ is n -integral over the ring $(A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]) [v]$. This proves Lemma 5.9 (b). \square

Proof of Theorem 5.8. (a) Since $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , Lemma 2.18 (applied to $A' = A[x]$) yields that $(I_\rho A[x])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[x]$.

Since $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , Lemma 2.18 (applied to $A' = A[y]$) yields that $(I_\rho A[y])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[y]$.

Since $(I_\rho)_{\rho \in \mathbb{N}}$ is an ideal semifiltration of A , Lemma 2.18 (applied to $A' = A[xy]$) yields that $(I_\rho A[xy])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[xy]$.

Thus, Theorem 5.8 (a) is proven.

(b) For every $v \in B$, the family $(I_\rho A[v])_{\rho \in \mathbb{N}}$ is an ideal semifiltration of $A[v]$ (by Lemma 2.18, applied to $A' = A[v]$), and thus we can consider the polynomial ring $(A[v])[Y]$ and its $A[v]$ -subalgebra $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right]$. For every $v \in B$, the polynomial ring $B[Y]$ is an $(A[v])[Y]$ -algebra (as explained in Definition 2.6), since B is an $A[v]$ -algebra¹³. Hence, this ring $B[Y]$ is an $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right]$ -algebra as well (because $(A[v]) \left[(I_\rho A[v])_{\rho \in \mathbb{N}} * Y \right]$ is a subring of $(A[v])[Y]$). Similarly, the ring $B[Y]$ is an $A \left[(I_\rho)_{\rho \in \mathbb{N}} * Y \right]$ -algebra.

¹³because $A[v]$ is a subring of B

Lemma 5.9 **(b)** (applied to $v = x$) yields that the element u of B is n -integral over $\left(A[x], (I_\rho A[x])_{\rho \in \mathbb{N}}\right)$ if and only if the element uY of the polynomial ring $B[Y]$ is n -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][x]\right)$. But since the element u of B is n -integral over $\left(A[x], (I_\rho A[x])_{\rho \in \mathbb{N}}\right)$, this yields that the element uY of the polynomial ring $B[Y]$ is n -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][x]\right)$.

Lemma 5.9 **(b)** (applied to y and m instead of v and n) yields that the element u of B is m -integral over $\left(A[y], (I_\rho A[y])_{\rho \in \mathbb{N}}\right)$ if and only if the element uY of the polynomial ring $B[Y]$ is m -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][y]\right)$. But since the element u of B is m -integral over $\left(A[y], (I_\rho A[y])_{\rho \in \mathbb{N}}\right)$, this yields that the element uY of the polynomial ring $B[Y]$ is m -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][y]\right)$.

Thus we know that uY is n -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][x]\right)$, and that uY is m -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][y]\right)$. Hence, Theorem 5.7 (applied to $A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right]$, $B[Y]$ and uY instead of A , B and u) yields that there exists some $\lambda \in \mathbb{N}$ such that uY is λ -integral over $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][xy]\right)$. Consider this λ .

Lemma 5.9 **(b)** (applied to xy and λ instead of v and n) yields that the element u of B is λ -integral over $\left(A[xy], (I_\rho A[xy])_{\rho \in \mathbb{N}}\right)$ if and only if the element uY of the polynomial ring $B[Y]$ is λ -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][xy]\right)$. But since the element uY of the polynomial ring $B[Y]$ is λ -integral over the ring $\left(A\left[(I_\rho)_{\rho \in \mathbb{N}} * Y\right][xy]\right)$, this yields that the element u of B is λ -integral over $\left(A[xy], (I_\rho A[xy])_{\rho \in \mathbb{N}}\right)$. Thus, Theorem 5.8 **(b)** is proven. \square

5.4. Second proof of Corollary 1.12

We notice that Corollary 1.12 can be derived from Lemma 5.1:

Second proof of Corollary 1.12. Let $n = 1$. Let $m = 1$. From $n = 1$, we obtain $\langle u^0, u^1, \dots, u^{n-1} \rangle_A = \langle u^0, u^1, \dots, u^0 \rangle_A = \langle u^0 \rangle_A = \langle 1_B \rangle_A$ (since $u^0 = 1_B$). Similarly, from $m = 1$, we obtain $\langle u^0, u^1, \dots, u^{m-1} \rangle_A = \langle 1_B \rangle_A$.

Now, we have

$$u^n \in \langle u^0, u^1, \dots, u^{n-1} \rangle_A \cdot \langle v^0, v^1, \dots, v^\alpha \rangle_A$$

¹⁴ and

$$u^m v^\beta \in \left\langle u^0, u^1, \dots, u^{m-1} \right\rangle_A \cdot \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A \\ + \left\langle u^0, u^1, \dots, u^m \right\rangle_A \cdot \left\langle v^0, v^1, \dots, v^{\beta-1} \right\rangle_A$$

¹⁵. Thus, Lemma 5.1 (applied to v , β and α instead of x , μ and ν) yields that u is $(n\beta + m\alpha)$ -integral over A (since $\beta + \alpha = \alpha + \beta \in \mathbb{N}^+$). This means that u is $(\alpha + \beta)$ -integral over A (because $\underbrace{n}_{=1}\beta + \underbrace{m}_{=1}\alpha = 1\beta + 1\alpha = \beta + \alpha = \alpha + \beta$).

This proves Corollary 1.12 once again. \square

¹⁴Proof. From $n = 1$, we obtain

$$u^n = u^1 = u = \sum_{\substack{i=0 \\ \in A}}^{\alpha} s_i v^i \in \left\langle v^0, v^1, \dots, v^\alpha \right\rangle_A = \left\langle u^0, u^1, \dots, u^{n-1} \right\rangle_A \cdot \left\langle v^0, v^1, \dots, v^\alpha \right\rangle_A,$$

since

$$\underbrace{\left\langle u^0, u^1, \dots, u^{n-1} \right\rangle_A}_{=\langle 1_B \rangle_A} \cdot \left\langle v^0, v^1, \dots, v^\alpha \right\rangle_A = \langle 1_B \rangle_A \cdot \left\langle v^0, v^1, \dots, v^\alpha \right\rangle_A = \left\langle 1_B v^0, 1_B v^1, \dots, 1_B v^\alpha \right\rangle_A \\ = \left\langle v^0, v^1, \dots, v^\alpha \right\rangle_A.$$

¹⁵Proof. We have

$$\underbrace{\left\langle u^0, u^1, \dots, u^{m-1} \right\rangle_A}_{=\langle 1_B \rangle_A} \cdot \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A = \langle 1_B \rangle_A \cdot \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A = \left\langle 1_B v^0, 1_B v^1, \dots, 1_B v^\beta \right\rangle_A \\ = \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A. \quad (54)$$

From $m = 1$, we obtain $u^m = u^1 = u$ and thus

$$\underbrace{u^m}_{=u} v^\beta = u v^\beta = \sum_{i=0}^{\beta} t_i v^{\beta-i} = \sum_{i=0}^{\beta} t_{\beta-i} \underbrace{v^{\beta-(\beta-i)}}_{=v^i}$$

(here we substituted $\beta - i$ for i in the sum)

$$= \sum_{\substack{i=0 \\ \in A}}^{\beta} t_{\beta-i} v^i \in \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A \\ = \left\langle u^0, u^1, \dots, u^{m-1} \right\rangle_A \cdot \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A \quad (\text{by (54)}) \\ \subseteq \left\langle u^0, u^1, \dots, u^{m-1} \right\rangle_A \cdot \left\langle v^0, v^1, \dots, v^\beta \right\rangle_A + \left\langle u^0, u^1, \dots, u^m \right\rangle_A \cdot \left\langle v^0, v^1, \dots, v^{\beta-1} \right\rangle_A.$$

References

- [1] Allen Altman, Steven Kleiman, *A Term of Commutative Algebra*, 1 September 2013.
<https://web.mit.edu/18.705/www/13Ed.pdf>
- [2] Nicolas Bourbaki, *Elements of Mathematics: Commutative Algebra*, Hermann 1972.
- [3] Antoine Chambert-Loir, *(Mostly) Commutative Algebra*, October 24, 2014.
<https://webusers.imj-prg.fr/~antoine.chambert-loir/enseignement/2014-15/ga/commalg.pdf>
- [4] J. S. Milne, *Algebraic Number Theory*, version 3.07.
<https://www.jmilne.org/math/CourseNotes/ant.html>
- [5] Craig Huneke and Irena Swanson, *Integral Closure of Ideals, Rings, and Modules*, London Mathematical Society Lecture Note Series, 336. Cambridge University Press, Cambridge, 2006.
<https://people.reed.edu/~iswanson/book/index.html>
- [6] Henri Lombardi, *Hidden constructions in abstract algebra (1) Integral dependance relations*, Journal of Pure and Applied Algebra 167 (2002), pp. 259–267.
<http://hlombardi.free.fr/publis/IntegralDependance.ps>
- [7] Darij Grinberg, *Integrality over ideal semifiltrations*, brief version.
<https://www.cip.ifi.lmu.de/~grinberg/algebra/integrality-merged.pdf>
- [8] Darij Grinberg, *Notes on the combinatorial fundamentals of algebra*, 10 January 2019.
<https://www.cip.ifi.lmu.de/~grinberg/primes2015/sols.pdf>
The numbering of theorems and formulas in this link might shift when the project gets updated; for a “frozen” version whose numbering is guaranteed to match that in the citations above, see <https://github.com/darijgr/detnotes/releases/tag/2019-01-10>.